TABLE S1. Antifungal susceptibility test

Drugs	MIC (μg/mL)														Range (μg/mL)	MIC 50	MIC 90	GM[a]	Mode[b]
	256	128	64	32	16	8	4	2	1	0.5	0.25	0.125	0.06	0.03					
AND	-	-	-	-	-	-	-	-	2	16	7	-	-	-	0.25-1	0.5	0.5	0.44	0.5
MF	-	-	-	-	-	-	-	2	17	6	-	-	-	-	0.5-2	1	1	0.90	1
CAS	-	-	-	-	-	-	-	-	9	15	1	-	-	-	0.25-1	0.5	1	0.62	0.5
ISA	-	-	-	-	-	-	-	-	-	-	5	10	7	3	0.03-0.25	0.125	0.125	0.09	0.125
PZ	-	-	-	-	-	-	-	-	-	-	-	8	11	6	0.03-0.125	0.06	0.125	0.06	0.06
VOR	-	-	-	-	-	-	1	19	5	-	-	-	-	-	1-4	2	2	1.79	2

IZ	-	-	-	-	-	-	-	-	-	-	5	19	1	-	0.06-0.25	0.125	0.125	0.13	0.125
FLZ	13	12	-	-	-	-	-	-	-	-	-	-	-	-	128-256	256	256	183.55	256
AB	-	-	-	-	-	-	-	2	23	-	-	-	-	-	1-2	1	1	1.06	1

Anidulafungin (AND), micafungin (MF), caspofungin (CAS), isavuconazole (ISA), posaconazole (PZ), voriconazole (VOR), itraconazole (ITZ), fluconazole (FLZ), and amphotericin B (AB). [a] Geometric mean calculated by multiplying all the MIC values and taking the n-th root, being n the number of values in the dataset [b] Mode calculated as the most frequently appeared MIC value in the dataset.