

Supplemental File S1.




Figure S1. Phase-contrast microscopy images of *C. auris* cells in RPMI-1640 with isolate IFRC2087 at 30 °C after 24 hours incubation. Magnification, $\times 200$.




Figure S2. Phase-contrast microscopy images of *C. auris* cells in RPMI-1640 with isolate IFRC2087 at 37 °C after 24 hours incubation. Magnification, $\times 200$.




Figure S3. Phase-contrast microscopy images of *C. auris* cells in RPMI-1640 with isolate IFRC4050 at 30 °C after 24 hours incubation. Magnification, $\times 200$.




Figure S4. Phase-contrast microscopy images of *C. auris* cells in RPMI-1640 with isolate IFRC4050 at 37 °C after 24 hours incubation. Magnification, $\times 200$.




Figure S5. Phase-contrast microscopy images of *C. auris* cells in RPMI-1640 with isolate MRL40 at 30 °C after 24 hours incubation. Magnification, $\times 200$.



Figure S6. Phase-contrast microscopy images of *C. auris* cells in RPMI-1640 with isolate MRL40 at 37 °C after 24 hours incubation. Magnification, $\times 200$.




Figure S7. Phase-contrast microscopy images of *C. auris* cells in RPMI-1640 with isolate TMML616 at 30 °C after 24 hours incubation. Magnification, $\times 200$.




Figure S8. Phase-contrast microscopy images of *C. auris* cells in RPMI-1640 with isolate TMML616 at 37 °C after 24 hours incubation. Magnification, $\times 200$.




Figure S9. Phase-contrast microscopy images of *C. auris* cells in RPMI-1640 with isolate TMML617 at 30 °C after 24 hours incubation. Magnification, $\times 200$.




Figure S10. Phase-contrast microscopy images of *C. auris* cells in RPMI-1640 with isolate TMML617 at 37 °C after 24 hours incubation. Magnification, $\times 200$.