

Supplementary Materials

***Beauveria bassiana* (Hypocreales: Clavicipitaceae) Volatile Organic Compounds (VOCs) Repel *Rhynchophorus ferrugineus* (Coleoptera: Dryophthoridae)**

Johari Jalinas ^{1,2,3}, Federico Lopez-Moya ^{1,*}, Frutos C. Marhuenda-Egea ^{2,4} and Luis Vicente Lopez-Llorca ^{1,2}

¹ Laboratory of Plant Pathology, Department of Marine Sciences and Applied Biology, University of Alicante, 03080 Alicante, Spain; johari_j@ukm.edu.my (J.J.); lv.lopez@ua.es (L.V.L.-L.)

² Multidisciplinary Institute for Environmental Studies (MIES) Ramon Margalef, University of Alicante, 03080 Alicante, Spain; frutos@gcloud.ua.es

³ Department of Biological Sciences and Biotechnology, Faculty of Science and Technology, Universiti Kebangsaan Malaysia (UKM), Selangor 43600, Malaysia

⁴ Department of Agrochemistry and Biochemistry, University of Alicante, 03080 Alicante, Spain

* Correspondence: federico.lopez@ua.es




Figure S1. Tube Y olfactometer to test the response of adult red palm weevil (female) on different chemical stimuli.




Figure S2. Response of a *Rhynchophorus ferrugineus* female in a Y-tube olfactometer (10 min) when given a choice between environmental air (control) and odour stimulus (Chemical stimuli) individuals per choice test. * Indicates significant difference within a choice test ($P < 0.01$, chi-square).