

Figure S1: Agarose gel of PCR products generated by using Cfl1\_for and Cfl1\_rev specific primers on DNA of clones obtained by ATCC20509 transformation using cfl1PT\_hy plasmid. M: marker, ctr+: cfl1PT\_hy plasmid.




Figure S2: Agarose gel of PCR products generated by using Cfl1\_for and Cfl1\_rev specific primers on cDNA (synthesized from mRNA) of clones obtained by ATCC20509 transformation using cfl1PT\_hy plasmid. M: marker, ctr+: cfl1PT\_hy plasmid.




Table S1: Primer sequences employed for cfl1PT\_hy assembly and mutant screening.

Primer	Sequence 5'-3'
hygr2_for	ATGAAGAAGCCCGAGCTCACCG
hygr2_rev	CGCTATCCGCGAGGACCTCG
Cfl1PT_HindIII_rev	tattAAGCTTCGCCCTCGATCAAGC
Cfl1PT_BcuI_for	atgtACTAGTTTGCGGCCGCTTTTCGAC
cfl1_for	gctcgttaccaacatctttgc
cfl1_rev	cactgggtgccattgagacg
SEQ3	CGCTCAAGTACTCCCTG
SEQ4	GGTTGGTGGTATTCCGC
SEQ5	GGCAGATGATGTCGAGG
CECK2	CCGAAGGTGTCGTTGGGTGCGCCG