

Table S2. *In vitro* antifungal susceptibility profile of 10 yeast species isolated from dogs (n=87) against different antifungal drugs using CLSI-BMD method.

Species (no. of strains tested)	Parameters	MICs (mg/L) ^a								
		FLU	ITC	VRC	ISA	POS	AMB	MFG	AFG	5-FC
<i>C. auris</i> (n=7)	Range	32->128	0.06-1	0.03-0.5	0.12-0.25	0.015-0.5	0.25-0.5	0.25-1	0.01-0.06	0.125-0.5
	MIC ₅₀ ^b	64	0.06	0.03	0.125	0.015	0.5	0.5	0.03	0.125
	MIC ₉₀ ^c	>128	0.7	0.5	0.75	0.35	0.5	1	0.06	0.35
<i>M. pachydermatis</i> (n=15)	Range	1-4	0.03-0.5	0.03-0.06	0.01-0.12	0.25-0.5	0.5	1	0.5-1	0.06
	MIC ₅₀	2	0.04	0.03	0.06	0.25	0.5	1	0.75	0.06
	MIC ₉₀	3.4	0.36	0.05	0.10	0.42	0.5	1	1	0.06
<i>C. tropicalis</i> (n=17)	Range	1-2	0.06-0.5	0.03-0.25	0.01-0.12	0.06-0.5	0.03-2	0.03-0.12	0.01	0.06-0.25
	MIC ₅₀	1	0.12	0.06	0.12	0.25	1	0.03	0.01	0.06
	MIC ₉₀	2	0.5	0.06	0.12	0.45	2	0.06	0.01	0.22
<i>C. krusei</i> (n=8)	Range	16-32	0.25-1	0.12-0.5	0.03-0.5	0.5	0.03-1	0.12-0.5	0.01-0.12	1-16
	MIC ₅₀	32	1	0.25	0.25	0.5	0.25	0.5	0.01	16
	MIC ₉₀	32	1	0.35	0.35	0.5	0.7	0.5	0.08	16
<i>T. asahii</i> (n=7)	Range	0.25-1	0.03-0.06	0.03-0.06	0.01-0.012	0.25-0.5	1-8	2-8	8	2-8
	MIC ₅₀	0.5	0.03	0.03	0.06	0.25	2.5	8	8	3
	MIC ₉₀	0.75	0.06	0.045	0.09	0.37	8	8	8	6
<i>C. lusitaniae</i> (n=4)	Range	1-2	0.12-1	0.03-0.06	0.06-0.25	0.25	0.5-1	0.12-0.25	0.01-0.06	0.06-0.12

	MIC ₅₀	1.5	0.56	0.04	0.15	0.25	0.75	0.18	0.03	0.09
	MIC ₉₀	2	1	0.06	0.25	0.25	1	0.25	0.06	0.12
<i>C. glabrata</i> (n=6)	Range	2	0.5	0.5	1	1	0.5	1	0.5-1	0.06
	MIC ₅₀	2	0.5	0.5	1	1	0.5	1	0.75	0.06
	MIC ₉₀	2	0.5	0.5	1	1	0.5	1	1	0.06
<i>C. rugosa</i> (n=4)	Range	2	0.03-0.06	0.03-0.06	0.12-0.25	0.01-0.06	0.5-2	0.12-0.25	0.01-0.03	0.06-0.12
	MIC ₅₀	2	0.03	0.03	0.25	0.01	0.5	0.25	0.03	0.06
	MIC ₉₀	2	0.05	0.05	0.25	0.04	1.55	0.25	0.03	0.10
<i>C. albicans</i> (n=3)	MIC ₅₀	2	0.03	0.06	0.25	0.5	0.25	0.25	0.03	0.5
<i>C. parapsilosis</i> (n=1)	MIC	1	0.125	0.06	0.06	0.25	1	0.5	0.5	0.06

^aFLU, fluconazole; ITC, itraconazole; VRC, voriconazole; ISA, isavuconazole; POS, posaconazole; AMB, amphotericin B; MFG, micafungin; AFG, anidulafungin; 5FC, flucytosine. ^b MIC₅₀, MIC at which 50% of test isolates were inhibited. ^c MIC₉₀, MIC at which 90% of test isolates were inhibited.