

Supplementary Material


Mouthrinse Is a Novel Non-Invasive Sampling Method for the Diagnosis and Screening of Sjögren Syndrome

Kouta Yamashiro ¹, Tomofumi Hamada ^{1,2}, Kazuki Mori ¹, Keitaro Nishi ¹, Maya Nakamura ¹, Mahiro Beppu ¹, Akihiko Tanaka ¹, Hiroshi Hijioka ¹, Yoshiaki Kamikawa ¹ and Tsuyoshi Sugiura ¹


¹ Department of Maxillofacial Diagnostic and Surgical Science, Field of Oral and Maxillofacial Rehabilitation, Graduate School of Medical and Dental Science, Kagoshima University, Kagoshima, Japan

² Department of Oral & Maxillofacial Surgery, Hakuai Medical Cooperation, Sagara Hospital, Kagoshima, Japan

* Correspondence: sugiura@dent.kagoshima-u.ac.jp



Supplementary Figure S1. Comparison of normalized signal intensity of six microRNAs between patients with Sjögren's syndrome (SS) and controls (HC). There were significant differences in signal intensity for miR-3648 ($p < 0.05$, t tests). Signal intensity did not significantly differ for the other five miRNAs. *: $p < 0.05$, ns: no significant difference.



Supplementary Figure S2. AUC of six microRNAs. The six microRNA's area under ROC curves (AUC) did not significantly discriminate Sjögren's syndrome (SS) patients from controls. ns: no significant difference.