Table S1. Effects of two-week PCS supplementation on the activity of antioxidant enzymes and GST in the kidneys of rats treated for four weeks with clozapine, aripiprazole, or risperidone. Enzyme activities were determined spectrophotometrically. Data are expressed as mean ± SEM (n = 8 per group) in units per milligram of total protein. Abbreviations: C, control group; C + PCS, control + PCS supplementation; Clo, clozapine group; Clo + PCS, clozapine + PCS supplementation; Ari, aripiprazole group; Ari + PCS, aripiprazole + PCS supplementation; Ris, risperidone group; Ris + PCS, risperidone + PCS supplementation. Statistical significance was assessed by two-way ANOVA, with results described in the text.

	Control	**Control+PCS**		**Clo**	**Clo+PCS**		**Ari**	**Ari + PCS**		**Ris**	**Ris + PCS**
CuZnSOD	24.97 ± 1.87	26.55 ± 1.00		18.91 ± 0.99	24.16 ± 1.73		19.92 ± 0.68	24.34 ± 1.73		21.01±1.27	27.25 ± 1.71
MnSOD	1.61 ± 0.10	1.52 ± 0.08		1.04 ± 0.07	1.18 ± 0.08		1.13 ± 0.05	1.00 ± 0.11		1.07 ± 0.05	1.36 ± 0.14
CAT	259.57 ± 11.44	272.56 ± 15.75	Clo	300.45 ± 19.71	362.23 ± 22.38	Ari	279.75 ± 25.01	295.59 ± 38.50	Ris	224.56 ± 14.12	264.86 ± 20.73
GPx	178.96 ± 7.64	254.73 ± 10.12		157.79 ± 3.68	167.57 ± 7.54		246.41 ± 6.62	298.95 ± 11.28		237.26 ± 6.62	273.39 ± 16.74
GR	238.82 ± 6.37	175.21 ± 9.54		237.04 ± 9.64	241.59 ± 14.17		176.33 ± 4.67	182.53 ± 5.83		174.24 ± 4.17	177.13 ± 7.81
GST	200.23 ± 9.21	191.04 ± 9.37		206.12 ± 8.22	201.50 ± 10.31		196.34 ± 10.23	202.60 ± 6.51		185.37 ± 7.56	191.71 ± 14.77