

Thermosensitive Drug Delivery System SBA-15-PEI for Controlled Release of Nonsteroidal Anti-inflammatory Drug Diclofenac Sodium Salt: A Comparative Study

Lubos Zauska¹, Stefan Bova², Eva Benova¹, Jozef Bednarcik³, Matej Balaz⁴, Vladimir Zelenak¹, Virginie Hornebecq⁵ and Miroslav Almasi^{1,*}

¹ Department of Inorganic Chemistry, Faculty of Science, P. J. Šafárik University, Moyzesova 11, SK-041 01 Košice, Slovakia; lubos.zauska@student.upjs.sk (L.Z.); evapopjakova@gmail.com (E.B.); vladimir.zelenak@upjs.sk (V.Z.)

² BovaChem s.r.o, Garbiarska 1919/14, SK-048 01, Rožňava, Slovakia; stefan.bova@gmail.com

³ Institute of Experimental Physics, Slovak Academy of Sciences, Watsonova 47, SK-040 01 Košice, Slovakia; jozef.bednarcik@upjs.sk

⁴ Institute of Geotechnics, Slovak Academy of Sciences, Watsonova 45, SK-040 01 Košice, Slovakia; balazm@saske.sk

⁵ Aix-Marseille University, CNRS, MADIREL, F-133 97 Marseille, France; virginie.hornebecq@univ-amu.fr

* Correspondence: miroslav.almasi@upjs.sk; Tel.: +421 552-342-366




Figure S1. (a) UV spectra of diclofenac sodium methanolic solutions and (b) corresponding calibration curve.




Figure S2. Calibration curve of diclofenac sodium in saline solutions at (a) pH = 2 and (b) pH = 7.4.




Figure S3. IR spectrum of diclofenac sodium salt.




Figure S4. Thermogravimetric curve of diclofenac sodium salt.