

Supplementary Materials

Oxidation of cyclohexanone with peracids – a straight path to the synthesis of ϵ -caprolactone oligomers

Jakub Bińczak^{1,2}, **Anna Szelwicka**³, **Agnieszka Siewniak**³, **Krzysztof Dziuba**¹ and **Anna Chrobok**^{3,*}

¹ Grupa Azoty Zakłady Azotowe „Puławy” S.A., Al. Tysiąclecia Państwa Polskiego 13, 24-110 Puławy, Poland

² Department of Chemical Organic Technology and Petrochemistry, PhD School, Silesian University of Technology, Akademicka 2A, 44-100 Gliwice, Poland

³ Department of Chemical Organic Technology and Petrochemistry, Faculty of Chemistry, Silesian University of Technology, Krzywoustego 4, 44-100 Gliwice, Poland

* Correspondence: anna.chrobok@polsl.pl; Tel.: +48 32 2371032




Figure S1. MS MALDI TOF of OCL isolated from reaction carried out in cyclohexane.




Figure S2. ^1H NMR of OCL isolated from reaction carried out in cyclohexane.




Figure S3. ¹H NMR analysis of OCL obtained in the reaction with CNON:perC₁₀ molar ratio 1:1.




Figure S4. ¹H NMR analysis of OCL obtained in the reaction with CNON:perC₁₀ molar ratio 1:2.




Figure S5. MS MALDI TOF analysis of OCL obtained in the reaction with CNON:perC₁₀ molar ratio 1:1.




Figure S6. MS MALDI TOF analysis of OCL obtained in the reaction with CNON:perC₁₀ molar ratio 1:2.