

Local Mo⁶⁺ Coordination and Textural Accessibility, Rather than Bulk Phase, Correlate with the Peroxidase-Mimetic Reactivity of CoMoO₄ Nanopowders

Milena Rosić ¹, Zorica Mojović ², Maja C. Pagnacco ², Jelena V. Senćanski ³, Jelena P. Maksimović ⁴, Dragana Jordanov ¹ and Maria Čebela ^{1,*}

¹ “Vinča” Institute of Nuclear Sciences, National Institute of the Republic of Serbia, University of Belgrade, Mike Petrovića Alasa 12-14, 11351 Belgrade, Serbia; mrosic@vin.bg.ac.rs (M.R.); djordano@vin.bg.ac.rs (D.J.)

² Institute of Chemistry, Technology and Metallurgy, University of Belgrade, Njegoševa 12, 11000 Belgrade, Serbia; zorica.mojovic@ihtm.bg.ac.rs (Z.M.); maja.pagnacco@ihtm.bg.ac.rs (M.C.P.)

³ Institute for Multidisciplinary Research, University of Belgrade, Kneza Višeslava 1, 11030 Belgrade, Serbia; jelenasencanski@gmail.com

⁴ Faculty of Physical Chemistry, University of Belgrade, Studentski trg 12-16, 11158 Belgrade, Serbia; jelena.maksimovic@ffh.bg.ac.rs

* Correspondence: mcebela@vin.bg.ac.rs

Supporting Information




Figure S1. Cyclic voltammograms recorded in the a) MA- HClO_4 ; b) MnSO_4 - HClO_4 ; c) KIO_3 - HClO_4 , and d) H_2O_2 - HClO_4 electrolytes without (dashed) and with (solid) the added CoMoO_4 at 450°C .




Figure S2. Cyclic voltammograms recorded in the a) MA- HClO_4 ; b) MnSO_4 - HClO_4 ; c) KIO_3 - HClO_4 ; and d) H_2O_2 - HClO_4 electrolytes without (dashed) and with (solid) the added CoMoO_4 at $1000\text{ }^\circ\text{C}$.