



Figure S1. Flowchart of data screening and inclusion of the participants of this study.

Table S1. Effective association rules when AoAC set to consequent

| Rule | Support | Confidence | Lift |
|---------------------------|---------|------------|-------|
| Age⇒AoAC | 49.04% | 87.40% | 1.030 |
| Hyperlipidemia⇒AoAC | 38.33% | 87.04% | 1.026 |
| TC⇒AoAC | 19.48% | 86.82% | 1.023 |
| LDL-C⇒AoAC | 26.44% | 86.21% | 1.016 |
| SBP⇒AoAC | 39.28% | 86.15% | 1.015 |
| Tooth decay⇒AoAC | 12.26% | 85.95% | 1.013 |
| Hypertension ⇒AoAC | 80.82% | 85.85% | 1.012 |
| Tooth loss or decay ⇒AoAC | 79.78% | 85.74% | 1.010 |
| Tooth loss ⇒AoAC | 76.91% | 85.66% | 1.009 |
| DM⇒AoAC | 20.74% | 85.38% | 1.006 |
| TG⇒AoAC | 12.37% | 85.14% | 1.003 |
| FPG⇒AoAC | 14.73% | 85.08% | 1.002 |

Support, Confidence and Lift of association rules are calculated based on the Apriori algorithm. “⇒” is used to indicate the direction of association rules.

SBP indicates systolic blood pressure; TC, total cholesterol; TG, triglyceride; LDL-C, low-density lipoprotein cholesterol; DM, diabetes mellitus; FPG, fasting plasma glucose.

Table S2. Baseline characteristics of participants according to presence of tooth loss

| | No tooth loss | Tooth loss | P value |
|-------------------------------------|----------------------|----------------------|---------|
| Number | 624 | 2006 | |
| AoAC score | 0.06 (0.06, 0.12) | 0.06 (0.06, 0.12) | 0.602 |
| Age, years | 69.70 ± 4.95 | 71.26 ± 5.26 | <0.001 |
| Female, n (%) | 343 (54.97) | 1129 (56.28) | 0.595 |
| SBP, mmHg | 135.83 ± 16.97 | 136.90 ± 17.58 | 0.183 |
| DBP, mmHg | 77.95 ± 10.03 | 76.83 ± 10.34 | 0.018 |
| BMI, kg/m ² | 24.08 ± 3.11 | 24.07 ± 3.17 | 0.929 |
| CHD, n (%) | 95 (15.22) | 251 (12.51) | 0.092 |
| Hypertension, n (%) | 444 (71.15) | 1509 (75.22) | 0.048 |
| Diabetes Mellitus n (%) | 156 (25.00) | 488 (24.33) | 0.773 |
| Hyperlipidemia, n (%) | 238 (38.14) | 703 (35.04) | 0.173 |
| WBC, 10 ⁹ /L | 6.30 (5.40, 7.40) | 6.30 (5.40, 7.30) | 0.88 |
| HGB, g/L | 134.86 ± 13.23 | 134.18 ± 13.51 | 0.272 |
| PLT, 10 ⁹ /L | 212.59 ± 50.73 | 206.76 ± 48.73 | 0.01 |
| TG, mmol/L | 1.33 (0.91, 1.81) | 1.26 (0.93, 1.73) | 0.267 |
| TC, mmol/L | 4.97 (4.35, 5.68) | 5.01 (4.32, 5.74) | 0.726 |
| LDL-C, mmol/L | 2.89 (2.31, 3.53) | 2.93 (2.29, 3.55) | 0.862 |
| HDL-C, mmol/L | 1.34 (1.15, 1.58) | 1.38 (1.16, 1.63) | 0.102 |
| FPG, mmol/L | 5.30 (4.75, 5.99) | 5.34 (4.80, 6.11) | 0.084 |
| eGFR, ml/min/1.73m ² | 67.44 (61.82, 74.02) | 66.87 (60.52, 73.04) | 0.031 |
| Smoking, n (%) | 82 (13.14) | 301 (15.00) | 0.277 |
| Drinking, n (%) | 135 (21.63) | 527 (26.27) | 0.023 |
| Physical activity hours/week, n (%) | | | 0.231 |
| <3h | 124 (19.87) | 459 (22.88) | |
| 3-10h | 435 (69.71) | 1328 (66.20) | |
| >10h | 65 (10.42) | 219 (10.92) | |

Continuous variables are summarized as mean ± standard deviation for normally distributed, and median (25th, 75th percentiles) for non-normally distributed. Categorical variables are summarized as count (percentage).

SBP indicates systolic blood pressure; DBP, diastolic blood pressure; BMI, body mass index; CHD, coronary heart disease; WBC, white blood cell count; HGB, hemoglobin; PLT, platelet; TC, total cholesterol; TG, triglyceride; LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol; FPG, fasting plasma glucose; eGFR estimated glomerular filtration rate.

Table S3. Baseline characteristics of participants according to presence of tooth decay

| | No tooth decay | Tooth decay | P value |
|-------------------------------------|----------------------|----------------------|---------|
| Number | 2371 | 259 | |
| AoAC score | 0.06 (0.06, 0.12) | 0.12 (0.06, 0.12) | 0.551 |
| Age, years | 70.89 ± 5.28 | 70.86 ± 4.79 | 0.912 |
| Female, n (%) | 1323 (55.80) | 149 (57.53) | 0.641 |
| SBP, mmHg | 136.43 ± 17.36 | 138.66 ± 18.09 | 0.05 |
| DBP, mmHg | 76.96 ± 10.16 | 78.33 ± 11.26 | 0.042 |
| BMI, kg/m ² | 24.08 ± 3.14 | 23.97 ± 3.30 | 0.589 |
| CHD, n (%) | 317 (13.37) | 29 (11.20) | 0.376 |
| Hypertension, n (%) | 1765 (74.44) | 188 (72.59) | 0.566 |
| Diabetes Mellitus n (%) | 589 (24.84) | 55 (21.24) | 0.228 |
| Hyperlipidemia, n (%) | 854 (36.02) | 87 (33.59) | 0.48 |
| WBC, 10 ⁹ /L | 6.30 (5.40, 7.40) | 6.20 (5.30, 7.20) | 0.439 |
| HGB, g/L | 134.30 ± 13.55 | 134.71 ± 12.45 | 0.641 |
| PLT, 10 ⁹ /L | 207.83 ± 49.36 | 211.05 ± 48.30 | 0.318 |
| TG, mmol/L | 1.28 (0.93, 1.78) | 1.21 (0.92, 1.62) | 0.177 |
| TC, mmol/L | 4.99 (4.32, 5.74) | 5.05 (4.34, 5.70) | 0.58 |
| LDL-C, mmol/L | 2.93 (2.29, 3.54) | 2.88 (2.36, 3.63) | 0.433 |
| HDL-C, mmol/L | 1.37 (1.16, 1.61) | 1.41 (1.19, 1.65) | 0.226 |
| FPG, mmol/L | 5.33 (4.79, 6.09) | 5.29 (4.78, 6.01) | 0.522 |
| eGFR, ml/min/1.73m ² | 66.95 (60.70, 73.23) | 67.20 (62.00, 73.33) | 0.43 |
| Smoking, n (%) | 338 (14.26) | 45 (17.37) | 0.208 |
| Drinking, n (%) | 578 (24.38) | 84 (32.43) | 0.006 |
| Physical activity hours/week, n (%) | | | 0.946 |
| <3h | 527 (22.23) | 56 (21.62) | |
| 3-10h | 1587 (66.93) | 176 (67.95) | |
| >10h | 257 (10.84) | 27 (10.42) | |

Continuous variables are summarized as mean ± standard deviation for normally distributed, and median (25th, 75th percentiles) for non-normally distributed. Categorical variables are summarized as count (percentage).

SBP indicates systolic blood pressure; DBP, diastolic blood pressure; BMI, body mass index; CHD, coronary heart disease; WBC, white blood cell count; HGB, hemoglobin; PLT, platelet; TC, total cholesterol; TG, triglyceride; LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol; FPG, fasting plasma glucose; eGFR estimated glomerular filtration rate.

Table S4. Baseline characteristics of participants according to presence of tooth loss or decay

| | No tooth loss or decay | Tooth loss or decay | P value |
|-------------------------------------|------------------------|----------------------|---------|
| Number | 567 | 2063 | |
| AoAC score | 0.06 (0.06, 0.12) | 0.06 (0.06, 0.12) | 0.826 |
| Age, years | 69.66 ± 5.05 | 71.23 ± 5.23 | <0.001 |
| Female, n (%) | 313 (55.20) | 1159 (56.18) | 0.713 |
| SBP, mmHg | 135.94 ± 16.94 | 136.84 ± 17.57 | 0.273 |
| DBP, mmHg | 77.93 ± 10.04 | 76.87 ± 10.33 | 0.03 |
| BMI, kg/m ² | 24.14 ± 3.11 | 24.05 ± 3.16 | 0.584 |
| CHD, n (%) | 91 (16.05) | 255 (12.36) | 0.026 |
| Hypertension, n (%) | 406 (71.60) | 1547 (74.99) | 0.115 |
| Diabetes Mellitus n (%) | 146 (25.75) | 498 (24.14) | 0.463 |
| Hyperlipidemia, n (%) | 218 (38.45) | 723 (35.05) | 0.148 |
| WBC, 10 ⁹ /L | 6.30 (5.40, 7.40) | 6.30 (5.40, 7.30) | 0.912 |
| HGB, g/L | 134.90 ± 13.30 | 134.19 ± 13.49 | 0.264 |
| PLT, 10 ⁹ /L | 212.84 ± 51.30 | 206.86 ± 48.62 | 0.01 |
| TG, mmol/L | 1.33 (0.91, 1.83) | 1.26 (0.93, 1.73) | 0.322 |
| TC, mmol/L | 4.98 (4.39, 5.71) | 5.00 (4.30, 5.74) | 0.681 |
| LDL-C, mmol/L | 2.94 (2.36, 3.54) | 2.92 (2.29, 3.55) | 0.648 |
| HDL-C, mmol/L | 1.35 (1.15, 1.58) | 1.38 (1.16, 1.63) | 0.208 |
| FPG, mmol/L | 5.32 (4.76, 6.02) | 5.33 (4.79, 6.10) | 0.34 |
| eGFR, ml/min/1.73m ² | 67.66 (61.82, 74.02) | 66.81 (60.53, 73.06) | 0.038 |
| Smoking, n (%) | 69 (12.17) | 314 (15.22) | 0.079 |
| Drinking, n (%) | 121 (21.34) | 541 (26.22) | 0.02 |
| Physical activity hours/week, n (%) | | | 0.097 |
| <3h | 109 (19.22) | 474 (22.98) | |
| 3-10h | 401 (70.72) | 1362 (66.02) | |
| >10h | 57 (10.05) | 227 (11.00) | |

Continuous variables are summarized as mean ± standard deviation for normally distributed, and median (25th, 75th percentiles) for non-normally distributed. Categorical variables are summarized as count (percentage).

SBP indicates systolic blood pressure; DBP, diastolic blood pressure; BMI, body mass index; CHD, coronary heart disease; WBC, white blood cell count; HGB, hemoglobin; PLT, platelet; TC, total cholesterol; TG, triglyceride; LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol; FPG, fasting plasma glucose; eGFR estimated glomerular filtration rate.

Table S5. Baseline characteristics of participants according to presence of tooth loss or decay

| Variables | HR | 95% CI | P value |
|---------------------|-------|-------------|---------|
| Model 1 | | | |
| Tooth loss | 1.377 | 1.221-1.553 | <0.001 |
| Tooth loss or decay | 1.458 | 1.286-1.652 | <0.001 |
| Model 2 | | | |
| Tooth loss | 1.352 | 1.199-1.525 | <0.001 |
| Tooth loss or decay | 1.433 | 1.265-1.625 | <0.001 |
| Model 3 | | | |
| Tooth loss | 1.356 | 1.202-1.530 | <0.001 |
| Tooth loss or decay | 1.437 | 1.267-1.629 | <0.001 |

Model 1: adjusted with age, gender, BMI, SBP, DBP, smoking, drinking.

Model 2: further adjusted with hypertension, hyperlipidemia, diabetes mellitus

Model 3: further adjusted with TG, LDL-C, HDL-C, FPG, eGFR.

HR indicates hazard ratio; CI, confidence interval; BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; TG, triglyceride; LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein; FPG, fasting plasma glucose; eGFR, estimated glomerular filtration rate.