

Morphology Controlled WO₃ for the Photocatalytic Oxidation of Methane to Methanol in Mild Conditions

Dumindu Premachandra^{1,2}, and Michael D. Heagy^{2*}

¹Department of Chemistry, New Mexico Institute of Mining and Technology
801, Leroy Pl, Socorro, NM, 87801 USA


²School of Chemistry and Materials Science, Rochester Institute of Technology
One Lomb Memorial Drive, Rochester, NY 14623 USA


*Corresponding author: mdhsch@rit.edu

Supplementary information

Table of Contents

1. Figure S1. PXRD spectra of WO₃ morphologies.....S3
2. Figure S2. WO₃ Flowers with different electron scavengers.....S3
3. Figure S3. The yield of methanol in the photocatalytic partial oxidation of methane for WO₃ flowers/2 mM H₂O₂S4
4. Figure S4. FESEM image of WO₃ nanowires with high agglomerationS4








Figure S1. PXRD spectra of WO₃ morphologies.




Figure S2. WO₃ Flowers with different electron scavengers (Reaction conditions; temperature = 50 °C, pH =~ 7, UV light irradiation, CH₄/N₂ bubbling with 100 mL/min).




Figure S3. The yield of methanol in the photocatalytic partial oxidation of methane for WO₃ flowers/2 mM H₂O₂ (Reaction conditions; temperature = 50 °C, pH = ~ 7, UV light irradiation, CH₄/N₂ bubbling with 100 mL/min).




Figure S4. FESEM image of WO₃ nanowires with high agglomeration