

Supplementary Materials

Three-Dimensional Kidney-on-a-Chip Assessment of Contrast-induced Kidney Injury: Osmolality and Viscosity

Kipro Kim¹, Beomgyun Jeong², Yun-Mi Lee³, Hyung-Eun Son³, Ji-Young Ryu³, Seokwoo Park^{3,4}, Jong Cheol Jeong³, Ho Jun Chin^{3,5}, Sejoong Kim^{3,5,*}

¹ Division of Nephrology and Hypertension, Department of Internal Medicine, Inha University College of Medicine, Incheon 22332, Korea; kpkidney@inha.ac.kr

² Research Center for Materials Analysis, Korea Basic Science Institute, Daejeon 34133, Korea; bjeong@kbsi.re.kr

³ Department of Internal Medicine, Seoul National University Bundang Hospital, Seongnam 13620, Korea; yunmi1202@hanmail.net (Y.-M.L.); she0817@naver.com (H.-E.S.); jyryu1022@gmail.com (J.-Y.R.); no1seokwoo@gmail.com (S.P.); jcj0425@empal.com (J.C.J.); mednep@snubh.org (H.J.C.)

⁴ Department of Biomedical Sciences, Seoul National University College of Medicine, Seoul 03080, Korea

⁵ Department of Internal Medicine, Seoul National University College of Medicine, Seoul 03080, Korea




Figure S1. Mathematical simulation of bidirectional flows in iopromide groups.




Figure S2. Mathematical simulation of bidirectional flows in iodixanol groups.