

Supplementary Materials

Trojan pH-Sensitive Polymer Particles Produced in a Continuous-Flow Capillary Microfluidic Device Using Water-in-Oil-in-Water Double-Emulsion Droplets

Ane Larrea ¹, Manuel Arruebo ^{1,2,3}, Christophe A. Serra ^{4,*} and Victor Sebastián ^{1,2,3,5,*}

¹ Instituto de Nanociencia y Materiales de Aragón (INMA), CSIC-Universidad de Zaragoza, 50009 Zaragoza, Spain; larrea@unizar.es (A.L.); arruebom@unizar.es (M.A.)


² Department of Chemical Engineering, Campus Río Ebro-Edificio I+D, University of Zaragoza, C/Poeta Mariano Esquillor S/N, 50018 Zaragoza, Spain

³ Networking Research Center on Bioengineering, Biomaterials and Nanomedicine, CIBER-BBN, 28029 Madrid, Spain

⁴ Université de Strasbourg, CNRS, ICS UPR 22, F-67000 Strasbourg, France

⁵ Laboratorio de Microscopías Avanzadas, Universidad de Zaragoza, 50018 Zaragoza, Spain

* Correspondence: ca.serra@unistra.fr (C.A.S.); victorse@unizar.es (V.S.)



Citation: Larrea, A.; Arruebo, M.; Serra, C.A.; Sebastián, V. Trojan pH-Sensitive Polymer Particles Produced in a Continuous-Flow Capillary Microfluidic Device Using Water-in-Oil-in-Water Double-Emulsion Droplets. *Micromachines* **2022**, *13*, 878. <https://doi.org/10.3390/mi13060878>

Academic Editor(s): Zebing Mao, Jin Xie and Hong Ding.

Received: date

Accepted: date

Published: 31 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure S2. Characterization of the enteric microparticles synthesized in the coaxial microfluidic system with different concentrations of Eudragit. $Q_i/Q_m/Q_o = 5/10/400 \mu\text{l}/\text{min}$. * CV is the ratio between the standard deviation and the average diameter.