

Supplementary material – Figure S1

Long time no hear, magnificent *Wohlfahrtia!* Morphological and molecular evidence of almost forgotten flesh fly in Serbia and Western Balkans

Stanislav Simin^{1,*}, Snežana Tomanović², Ratko Sukara², Marijana Stefanov¹, Milan Savović³, Bojan Gajić⁴ and Vesna Lalošević¹

¹University of Novi Sad, Faculty of Agriculture, Department of Veterinary Medicine, Trg Dositeja Obradovića 8, 21000 Novi Sad, Serbia; stanislav.simin@polj.edu.rs, marijana.stefanov96@gmail.com, lvesna@polj.uns.ac.rs

²University of Belgrade, Institute for Medical Research- National Institute of Republic of Serbia, Group for Medical Entomology, Centre of Excellence for Food- and Vector-Borne Zoonoses, 11129 Belgrade, Serbia; snezanat@imi.bg.ac.rs, ratko.sukara@imi.bg.ac.rs

³Private Veterinary Practice „MSV Medicus“, Milice Stojadinović Srpkinja 1, 21209 Bukovac, Serbia; msvmedicus@gmail.com

⁴United Arab Emirates University, College of Agriculture and Veterinary Medicine, Department of Veterinary Medicine, P.O. Box 15551, Al Ain, United Arab Emirates; b.gajic@uaeu.ac.ae

*Correspondence: stanislav.simin@polj.edu.rs




Figure S1. The location of a sheep farm where the cases of traumatic myiasis were recorded (the village of Jarkovac (45°16'05"N; 20°45'18"E), Central Banat District, Vojvodina Province, Northern Serbia).