






Supplementary Figure S1: Taxonomic profile (Class level) and relative abundance of the oral microbiota of the five groups of patients with different clinical condition of HIV. HP: Controls (without HIV), C_HIV: Virally suppressed, ND_HIV: Recently diagnosed HIV, Candida_HIV: HIV and oral candidiasis, HPV_HIV: HIV and HPV-related oral lesions.



Supplementary Figure S2: Taxonomic profile (Order level) and relative abundance of the oral microbiota of the five groups of patients with different clinical condition of HIV. HP: Controls (without HIV), C_HIV: Virally suppressed, ND_HIV: Recently diagnosed HIV, Candida_HIV: HIV and oral candidiasis, HPV_HIV: HIV and HPV-related oral lesions.



Supplementary Figure S3: Taxonomic profile (Family level) and relative abundance of the oral microbiota of the five groups of patients with different clinical condition of HIV. HP: Controls (without HIV), C_HIV: Virally suppressed, ND_HIV: Recently diagnosed HIV, Candida_HIV: HIV and oral candidiasis, HPV_HIV: HIV and HPV-related oral lesions.



Supplementary Figure S4: Taxonomic profile (Species level) and relative abundance of the oral microbiota of the five groups of patients with different clinical condition of HIV. HP: Controls (without HIV), C_HIV: Virally suppressed, ND_HIV: Recently diagnosed HIV, Candida_HIV: HIV and oral candidiasis, HPV_HIV: HIV and HPV-related oral lesions.