

# Secondary Metabolites, Including a New 5,6-Dihydropyran-2-one, Produced by the Fungus *Diplodia corticola*. Aphicidal Activity of the Main Metabolite, Sphaeropsidin A

Maria Michela Salvatore<sup>1,2a</sup>, Ilaria Di Lelio<sup>3a</sup>, Marina DellaGreca<sup>1</sup>, Rosario Nicoletti<sup>3,4</sup>, Francesco Salvatore<sup>1</sup>, Elia Russo<sup>3</sup>, Gennaro Volpe<sup>3</sup>, Andrea Becchimanz<sup>i3</sup>, Alla Eddine Mahamed<sup>i5</sup>, Akila Berraf-Tebbal<sup>6</sup>, and Anna Andolfi<sup>1,7\*</sup>

<sup>1</sup> Department of Chemical Sciences, University of Naples Federico II, 80126 Naples, Italy; mariamichela.salvatore@unina.it (M.M.S.); dellagre@unina.it (M.D.); frsalvat@unina.it (F.S.)

<sup>2</sup> Institute for Sustainable Plant Protection, National Research Council, 80055 Portici, Italy

<sup>3</sup> Department of Agriculture, University of Naples Federico II, 80055 Portici, Italy; ilaria.dilelio@unina.it (I.D.L.); rosario.nicoletti@crea.gov.it (R.N.); elia.russo@unina.it (E.R.); gennaro.volpe2@unina.it (G.V.); andrea.becchimanz<sup>i</sup>@unina.it (A.B.)

<sup>4</sup> Council for Agricultural Research and Economics, Research Center for Olive, Fruit, and Citrus Crops, 81100 Caserta, Italy

<sup>5</sup> Department of Biology, Faculty of Natural Sciences, Life and Earth Sciences, University of Ghardaia, 47000 Ghardaia, Algeria; aladin1342@yahoo.com

<sup>6</sup> Mendeleum-Institute of Genetics, Faculty of Horticulture, Mendel University in Brno, 69144 Lednice, Czech Republic; berraf.a@hotmail.fr

<sup>7</sup> BAT Center-Interuniversity Center for Studies on Bioinspired Agro-Environmental Technology, University of Naples Federico II, 80055 Portici, Italy

\* Correspondence: andolfi@unina.it

† These authors contributed equally to this work.

## Contents

**Figure S1.**  $^1\text{H}$  NMR spectrum of diplopyrone C (**1**) recorded in  $\text{CDCl}_3$  at 400 MHz

**Figure S2.**  $^{13}\text{C}$  NMR spectrum of diplopyrone (**1**) recorded in  $\text{CDCl}_3$  at 100 MHz

**Figure S3.**  $^1\text{H},^1\text{H}$  COSY spectrum of diplopyrone C (**1**) recorded in  $\text{CDCl}_3$  at 400 MHz

**Figure S4.** HSQC spectrum of diplopyrone C (**1**) recorded in  $\text{CDCl}_3$  at 400 MHz

**Figure S5.** HMBC spectrum of diplopyrone C (**1**) recorded in  $\text{CDCl}_3$  at 400 MHz

**Figure S6.** NOESY spectrum of diplopyrone C (**1**) recorded in  $\text{CDCl}_3$  at 400 MHz

**Figure S7.** IR spectrum of diplopyrone C (**1**)

**Figure S8.** HRESI MS spectrum of diplopyrone C (**1**) recorded in positive mode

**Figure S9.**  $^1\text{H}$  NMR spectrum of sphaeropsidin A (**2**) recorded in  $\text{CDCl}_3$  at 400 MHz

**Figure S10.**  $^1\text{H}$  NMR spectrum of sphaeropsidin B (**3**) recorded in  $\text{CDCl}_3$  at 400 MHz

**Figure S11.**  $^1\text{H}$  NMR spectrum of sphaeropsidin C (**4**) recorded in  $\text{CDCl}_3$  at 400 MHz

**Figure S12.**  $^1\text{H}$  NMR spectrum of (*3R*)-mellein (**5**) recorded at 400 MHz in  $\text{CDCl}_3$

**Figure S13.**  $^1\text{H}$  NMR spectrum of (*3R,4R*)-4-hydroxymellein (**6**) recorded at 400 MHz in  $\text{CDCl}_3$


**Figure S14.**  $^1\text{H}$  NMR spectrum of (*3R,4S*)-4-hydroxymellein (**7**) recorded at 400 MHz in  $\text{CDCl}_3$

**Figure S15.**  $^1\text{H}$  NMR spectrum sapinofuranone B (**8**) recorded at 400 MHz in  $\text{CDCl}_3$


**Figure S16.**  $^1\text{H}$  NMR spectrum of pinofuranoxin A (**9**) recorded at 400 MHz in  $\text{CDCl}_3$

**Figure S17.**  $^1\text{H}$  NMR spectrum of diplobifuranylone B (**10**) recorded in  $\text{CDCl}_3$  at 400 MHz


**Figure S18.**  $^1\text{H}$  NMR spectrum tyrosol (**11**) recorded at 400 MHz in  $\text{CDCl}_3$




**Figure S1.** <sup>1</sup>H NMR spectrum of diplopyrone C (1) recorded in CDCl<sub>3</sub> at 400 MHz.




**Figure S2.** <sup>13</sup>C NMR spectrum of diplopyrone (**1**) recorded in CDCl<sub>3</sub> at 100 MHz.




**Figure S3.** <sup>1</sup>H, <sup>1</sup>H COSY spectrum of diplopyrone C (**1**) recorded in CDCl<sub>3</sub> at 400 MHz.




**Figure S4.** HSQC spectrum of diplopyrone C (**1**) recorded in  $\text{CDCl}_3$  at 400 MHz.




**Figure S5.** HMBC spectrum of diplopyrone C (**1**) recorded in  $\text{CDCl}_3$  at 400 MHz.




**Figure S6.** NOESY spectrum of diplopyrone C (**1**) recorded in  $\text{CDCl}_3$  at 400 MHz.




**Figure S7.** IR spectrum of diplopyrone C (1).




**Figure S8.** HRESI MS spectrum of diplopyrone C (**1**) recorded in positive mode.




**Figure S9.** <sup>1</sup>H NMR spectrum of sphaeropsidin A (**2**) recorded in CDCl<sub>3</sub> at 400 MHz.




**Figure S10.** <sup>1</sup>H NMR spectrum of sphaeropsidin B (3) recorded in CDCl<sub>3</sub> at 400 MHz.




**Figure S11.** <sup>1</sup>H NMR spectrum of sphaeropsidin C (**4**) recorded in CDCl<sub>3</sub> at 400 MHz.




**Figure S12.** <sup>1</sup>H NMR spectrum of (3*R*)-mellein (**5**) recorded at 400 MHz in CDCl<sub>3</sub>.




**Figure S13.** <sup>1</sup>H NMR spectrum of (3*R*,4*R*)-4-hydroxymellein (**6**) recorded at 400 MHz in CDCl<sub>3</sub>.




**Figure S14.** <sup>1</sup>H NMR spectrum of (3*R*,4*S*)-4-hydroxymellein (**7**) recorded at 400 MHz in CDCl<sub>3</sub>.




**Figure S15.**  $^1\text{H}$  NMR spectrum sapinofuranone B (8) recorded at 400 MHz in  $\text{CDCl}_3$ .



**Figure S16.**  $^1\text{H}$  NMR spectrum of pinofuranoxin A (**9**) recorded at 400 MHz in  $\text{CDCl}_3$ .



**Figure S17.** <sup>1</sup>H NMR spectrum of diplobifuranylon B (**10**) recorded in CDCl<sub>3</sub> at 400 MHz.



**Figure S18.** <sup>1</sup>H NMR spectrum tyrosol (**11**) recorded at 400 MHz in CDCl<sub>3</sub>.