

Exploration of Cytotoxic Potential of Longifolene/Junipene Isolated from *Chrysopogon zizanioides*

Madhuri Grover ^{1,2}, Tapan Behl ^{3,*}, Tarun Virmani ², Mohit Sanduja ⁴, Hafiz A. Makeen ⁵, Mohammed Albratty ⁶, Hassan A. Alhazmi ^{6,7}, Abdulkarim M. Meraya ⁵ and Simona Gabriela Bungau ^{8,9,*}

¹ Bhawani Shankar (B.S.) Anangpuria Institute of Pharmacy, Alampur, Ballabgarh 121004, India

² School of Pharmaceutical Sciences, Modern vidya Niketan MVN University, Palwal 121105, Haryana, India

³ School of Health Sciences, University of Petroleum and Energy Studies,

Dehradun 248007, Uttarakhand, India

⁴ Department of Pharmacy, School of Medical and Allied Sciences, GD Goenka University, Gurugram 122103, Haryana, India

⁵ Pharmacy Practice Research Unit, Clinical Pharmacy Department, College of Pharmacy, Jazan University, Jazan 45142, Saudi Arabia

⁶ Department of Pharmaceutical Chemistry, College of Pharmacy, Jazan University, Jazan 45142, Saudi Arabia

⁷ Substance Abuse and Toxicology Research Centre, Jazan University, Jazan 45142, Saudi Arabia

⁸ Department of Pharmacy, Faculty of Medicine and Pharmacy, University of Oradea, 410087 Oradea, Romania

⁹ Doctoral School of Biomedical Sciences, University of Oradea, 410087 Oradea, Romania

* Correspondence: tapanbehl31@gmail.com (T.B.); simonabungau@gmail.com (S.G.B.)




Figure S1. HPLC analysis of isolated unknown compound 'X'.




Figure S2. FTIR of the standard and isolated unknown compound 'X'.




Figure S3. Proton [¹H] NMR of isolated unknown compound 'X'.




Figure S4. Carbon [^{13}C] NMR of isolated unknown compound 'X'.




Figure S5. LC-MS of isolated unknown compound 'X'.