

Supplementary Information

Contents:

Figure S1. Oxidation peak currents (I_{pa} , red) and I_{pa}/I_{pc} ratio (blue) on different electrodes for 20.0 $\mu\text{mol L}^{-1}$ rutin (scan rate: 0.1 V s^{-1}).

Figure S2. (a) Plots of $I - v$, and (b) plots of $E - \ln v$ of 20.0 $\mu\text{mol L}^{-1}$ rutin on NF/AuNPs/N-CPDs@FLBP/CILE at different scan rate (0.03, 0.05, 0.08, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7 and 0.8 V s^{-1}).

Figure S3. The oxidation peak current of (a) repeatability and (b) reproducibility for detection of 10.0 $\mu\text{mol L}^{-1}$ rutin.




Figure S1. Oxidation peak currents (I_{pa} , red) and I_{pa}/I_{pc} ratio (blue) on different electrodes for 20.0 $\mu\text{mol L}^{-1}$ rutin (scan rate: 0.1 V s^{-1}).




Figure S2. (a) Plots of $I - v$, and (b) plots of $E - \ln v$ of $20.0 \mu\text{mol L}^{-1}$ rutin on NF/AuNPs/N-CPDs@FLBP/CILE at different scan rate ($0.03, 0.05, 0.08, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7$ and 0.8 V s^{-1}).




Figure S3. The oxidation peak current of (a) repeatability and (b) reproducibility for detection of $10.0 \mu\text{mol L}^{-1}$ rutin.