Supplementary data

Anti-adipogenic Effects of Salicortin from *Salix pseudolasiogyne* Twigs in 3T3-L1 Cells

Hee Jung Kim [1,2,†], Da Eun Lee [3,†], Eon Chung Park [3], Moon-Jin Ra [4], Sang-Mi Jung [4], Jeong-Nam Yu [5], Sung Hee Um [1,2,6,*], and Ki Hyun Kim [3,*]

[1] Department of Molecular Cell Biology, Samsung Biomedical Research Institute, Sungkyunkwan University School of Medicine, Suwon, Gyeonggi-do, 16419, Republic of Korea
[2] Department of Health Sciences and Technology, Samsung Advanced Institute for Health Sciences and Technology, Samsung Medical Center, Sungkyunkwan University, Seoul, 06351, Republic of Korea
[3] School of Pharmacy, Sungkyunkwan University, Suwon 16419, Republic of Korea
[4] Hongcheon Institute of Medicinal Herb, Hongcheon-gun, Gangwon-do 25142, Republic of Korea
[5] Nakdonggang National Institute of Biological Resources, Sangju, Gyeongsangbuk-do 37242, Republic of Korea
[6] Biomedical Institute Convergence at Sungkyunkwan University, Suwon, Gyeonggi-do, 16419, Republic of Korea
* Correspondence: shum@skku.edu (S.H.U.); khkim83@skku.edu (K.H.K.); Tel.: +82-31-299-6123 (S.H.U.); +82-31-290-7700 (K.H.K.)
† These authors contributed equally to this study

Figure S1. ^{1}H NMR spectrum of **1** (CD_3OD, 850 MHz)




Figure S2. ^{1}H NMR spectrum of **2** (CD_3OD, 850 MHz)

