

Supporting Information

N-rich, polyphenolic porous organic polymer and its *invitro* anticancer activity on colorectal cancer

*Sabuj Kanti Das,[‡] Snehasis Mishra[‡], Krishna Das Saha, Debraj Chandra, Michikazu Hara, Amany A. Mostafa and Asim Bhaumik**

**Corresponding author. E-mail: msab@iacs.res.in*

[‡]These authors contributed equally.

Table of Contents

Figure S1 FTIR spectra of TrzTFPPOP after DI water, 2 (M) H₂SO₄ and 2 (M) NaOH treatments for 7 days.

Figure S2 Wide angle PXRD pattern of TrzTFPPOP.

Figure S3 MTT data of (0-10 µg/ml) TrzTFPPOP against HCT 116 cell line. The IC₃₀, IC₅₀ and IC₇₀ values are in the inset.

Figure S4 MTT data of (0-20 µg/ml) TrzTFPPOP against HEK293 cell line




Figure S1: FTIR spectra of TrzTFPPOP after DI water, 2 (M) H₂SO₄ and 2 (M)NaOH treatment for 7 days.




Figure S2: Wide angle PXRD pattern of TrzTFPPOP.




Figure S3: Percentage of cell death after treatment of (0-10 $\mu\text{g/ml}$) TrzTFPPOP. The IC₃₀, IC₅₀ and IC₇₀ values are in the inset.




Figure S4: Percentage of cell death after treatment of (0-10 µg/ml) TrzTFPPOP in HEK293 cell line.