

Supporting Information

Coumarins from Jinhua Finger Citron: Separation by Liquid–Liquid Chromatography and Potential Antitumor Activity

Chaoyue Wang ^{1,2,†}, Jiangang Huang ^{2,†}, Zhiling Zhou ¹, Ping Xu ³, Jingyi Shi ¹,
Yushun Yang ¹, Shengqiang Tong ^{3,*} and Hongyu Hu ^{2,*}

¹ Jinhua Advanced Research Institute, Jinhua 321015, China

² Xingzhi College, Zhejiang Normal University, Lanxi 321100, China

³ College of Pharmaceutical Science, Zhejiang University of Technology, Hangzhou 310014, China

* Correspondence: sqtong@zjut.edu.cn (S.T.); huhongyu22@126.com (H.H.)

† These authors contributed equally to this work.

Supplementary Figures

Representative NMR spectra:




Figure S1. The ¹H NMR spectrum of compound I.




Figure S2. The ¹³C NMR spectrum of compound I.




Figure S3. The ¹H NMR spectrum of compound II.




Figure S4. The ¹³C NMR spectrum of compound II.




Figure S5. The ¹H NMR spectrum of compound III.




Figure S6. The ¹³C NMR spectrum of compound III.




Figure S7. The ¹H NMR spectrum of compound IV.




Figure S8. The ¹³C NMR spectrum of compound IV.




Figure S9. The ¹H NMR spectrum of compound V.




Figure S10. The ^{13}C NMR spectrum of compound V.

Original Western Blotting images:

PARP



Bcl-2



GAPDH

