

Article

Phloretin Inhibits Quorum Sensing and Biofilm Formation in *Serratia marcescens*

Yueheng Qi ^{1,2,†}, Pengcheng Ji ^{3,†}, Kunyuan Yin ³, Yi Zheng ³, Jiangxiu Niu ², Aiqun Jia ¹, Jinwei Zhou ^{3,*} and Jingguo Li ^{1,*}

¹ Henan Provincial People's Hospital, People's Hospital of Zhengzhou University, Zhengzhou 477150, China;

² Luoyang Key Laboratory of Organic Functional Molecules, College of Food and Drug, Luoyang Normal University, Luoyang 471934, China;

³ School of Food and Biological Engineering, Xuzhou University of Technology, Xuzhou 221018, China;

* Correspondence: zhoujinwei@xzit.edu.cn (J.Z.); lijingguo@zzu.edu.cn (J.L.); Tel.: +86-516-83105383 (J.Z.); +86-371-65896979 (J.L.); Fax: +86-516-83105383 (J.L.)

[†] These authors contributed equally to this work.

Table S1 PCR primers for qRT-PCR.

Genes	Primer direction	Sequence (5'–3')
<i>katG</i>	F	GACTTCTTCCGCCACCTGTTCG
	R	CGATGTCTTTCGCCACCCACTG
<i>bsmA</i>	F	TAGTCCGCACACTCATCGC
	R	GATCTCCTGCGCCTGTGC
<i>bsmB</i>	F	GCGGATGTGTATGCCTTCG
	R	GCCACGCATTTCTTCACTCA
<i>fimC</i>	F	CCAAATCCACGCCGGAAC
	R	TTGTTCTTGTCGCTGTCCGT
<i>ebp</i>	F	CTCAGAGGCACACTGGGCATTG
	R	ATTGTCAGCACCGATGGCGATC
<i>pigM</i>	F	GTTTCGCTCAGTATCGCAGGTGAC
	R	GGTAGGCTGCCGTTTCGTTGAAG
<i>pigC</i>	F	TTCGTCACAAACCGCACTATT
	R	CGTCTTTCACCGCCCAT
<i>htpX</i>	F	GCATCCTGGCGAGCATCATCAC
	R	GCAAGGCGGCGATCATCTTCTC
<i>rplT</i>	F	ATGAAGCAGGCGAAAGGTTACTACG
	R	CGACGGTCACGGTAAGCATACTG