

SUPPLEMENTARY MATERIAL

Toxicokinetics and Tissue Distribution of the Hepatotoxic Triterpenoid Saponin Pterocephin A in Rats Using the Ultra-Performance Liquid Chromatography–Tandem Mass Spectrometry (UPLC-MS/MS) Method

Yiran Xiong ^{1,2}, Zhaoyue Dong ¹, Hongxu Zhou ^{1,3}, Jingxin Mao ², Lingjiang Zeng ⁴, Yunbin Jiang ¹, Fancheng Meng ¹, Zhihua Liao ⁴, Min Chen ^{1,2,*}

Contents of Supplementary Information.

Figure S1. ¹³C-NMR of PA (A), ¹H-NMR of PA (B), ¹³C-NMR of IS (C), ¹H-NMR of IS (D).

Figure S2. Representative chromatograms of PA and IS under methanol-0.1% ammonia.

Figure S3. Secondary mass spectra of PA under different crushing voltages.

Figure S4. Chromatograms for pre-treatment examination of blank plasma (A), PA containing plasma (B), blank tissue (C) and PA containing tissue (D).




Figure S1. ^{13}C -NMR of PA (A), ^1H -NMR of PA (B), ^{13}C -NMR of IS (C), ^1H -NMR of IS (D).




Figure S2. Representative chromatograms of PA and IS under methanol-0.1% ammonia.




Figure S3. Secondary mass spectra of PA under different crushing voltages.




Figure S4. Chromatograms for pre-treatment examination of blank plasma (A), PA containing plasma (B), blank tissue (C) and PA containing tissue (D).