SUPPLEMENTARY MATERIAL

Thermal Properties of Pharmaceutical Antioxidants: Heat Capacities and Melting Temperatures

Molecules

Iván Montenegro[a], Carmen Pérez[b], Begoña González[a], Ángeles Domínguez[a], Elena Gómez[a,*]

[a]FEQx lab, Department of Chemical Engineering, University of Vigo, Spain
[b]CINTECX, ENCOMAT Group, Material Science Department, CINTECX, University of Vigo, Spain

* Corresponding author
E-mail address: elenagc@uvigo.es

Figure Index

Figure S1 Chemical structure of *trans*-resveratrol, *trans*-polydatin, kaempferol, quercetin, myricetin, (-)-epicatechin, and hesperidin 2
Figure S2 DSC and TG curves of *trans*-polydatin 3
Figure S3 DSC and TG curves of kaempferol 3
Figure S4 DSC and TG curves of quercetin 4
Figure S5 DSC and TG curves of myricetin 4
Figure S6 DSC and TG curves of hesperidin 5
Figure S7 DSC and TG curves of (-)-epicatechin 5




Figure S1 Chemical structure of *trans*-resveratrol (a), *trans*-polydatin (b), kaempferol (c), quercetin (d), myricetin (e), (-)-epicatechin (f), and hesperidin (g)




Figure S2 DSC (—) and TG (---) curves of *trans*-polydatin




Figure S3 DSC (—) and TG (---) curves of kaempferol




Figure S4 DSC (—) and TG (---) curves of quercetin




Figure S5 DSC (—) and TG (---) curves of myricetin




Figure S6 DSC (—) and TG (---) curves of hesperidin




Figure S7 DSC (—) and TG (---) curves of (-)-epicatechin