

Supplementary information

pH-Sensitive Nanoparticles for Colonic Delivery Anti-miR-301a in Mice Models of Inflammatory Bowel Diseases

Junshan Wang ^{1,*}, Min Yao ², Jiafeng Zou ², Wenxing Ding ², Mingyue Sun ², Ying Zhuge ³ and Feng Gao ^{2,4,5,6,*}

¹ Department of Gastroenterology, Chongming Branch of Shanghai Tenth People's Hospital, Tongji University School of Medicine, Shanghai 202157, China

² Shanghai Frontier Science Research Base of Optogenetic Techniques for Cell Metabolism, School of Pharmacy, East China University of Science and Technology, Shanghai 200237, China; ym1584007995@163.com (M.Y.); zoujiafeng2019@126.com (J.Z.); dwx150120413@gmail.com (W.D.); 17854280576@163.com (M.S.)

³ Department of Cardiology, Shanghai General Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China; yiliaobangzu@163.com

⁴ Shanghai Key Laboratory of New Drug Design, School of Pharmacy, East China University of Science and Technology, Shanghai 200237, China

⁵ Shanghai Key Laboratory of Functional Materials Chemistry, East China University of Science and Technology, Shanghai 200237, China

⁶ Optogenetics and Synthetic Biology Interdisciplinary Research Center, State Key Laboratory of Bioreactor Engineering, East China University of Science and Technology, Shanghai 200237, China

* Correspondence: jswang22@126.com (J.W.); fgao@ecust.edu.cn (F.G.)

Supplementary Figures




Figure S1. The stability of anti-miR@HCeP NP at 1, 4 and 7 d.




Figure S2. The cell viability of FAM-anti-miR@HCeP NP