

Suppl. Table S2. Tick species removed from red deer, showing distribution of piroplasm-infected ticks. Infected ticks were PCR positive for either *Theileria* sp. OT3 or *T. capreoli*.

Red deer	tick	n	<i>Theileria</i> sp. OT3	<i>Theileria</i> <i>capreoli</i>
1	<i>I. ricinus</i>	10	-	1
2	<i>I. ricinus</i>	2	-	-
	<i>R. sanguineus</i> s.l.	1	-	-
3	<i>I. ricinus</i>	5	-	-
	<i>R. sanguineus</i> s.l.	2	-	-
4	<i>I. ricinus</i>	1	-	1
	<i>R. sanguineus</i> s.l.	4	-	-
	<i>D. marginatus</i>	1	-	-
5	<i>I. ricinus</i>	6	-	3
6	<i>I. ricinus</i>	1	-	-
	<i>R. sanguineus</i> s.l.	6	-	-
7	<i>I. ricinus</i>	2	-	-
	<i>R. sanguineus</i> s.l.	1	1	-
	<i>D. marginatus</i>	1	-	-
8	<i>R. sanguineus</i> s.l.	6	-	-
9	<i>I. ricinus</i>	2	-	-
10	<i>R. sanguineus</i> s.l.	3	-	1
11	<i>I. ricinus</i>	1	-	-
12	<i>I. ricinus</i>	1	-	-