

**Table S1.** Sequences of primers used for cytokine real-time PCR (qPCR).

Gene	Forward Primer	Reverse Primer	Reference
<i>beta-actin</i>	5'-GGCACCAACACCTTCTACAATG-3'	5'-TGGATGGCTACGTACATGGCTG-3'	45
<i>IFN-gamma</i>	5'-TGAAAATCCTGCAGAGCCAGAT-3'	5'-AGCTCATTGAATGCTTGGCG-3'	45
<i>IL4</i>	5'-TGACGGCACAGAGCTATTGATG-3'	5'-TTCTTCGTTGCTGTGAGGACG-3'	45
<i>IL10</i>	5'-AGGCCTGTCATCGATTCTC-3'	5'-TGGCCTGTAGACACCTTGGTC-3'	45
<i>TNF<math>\alpha</math></i>	5'-CTGTAGCCCACGTCGTAGC-3'	5'-TTGAGATCCATGCCGTTG-3'	46

## References

45. Varona, R.; Cadenas, V.; Gómez, L.; Martínez-A, C.; Márquez, G. CCR6 Regulates CD4+ T-cell-mediated acute graft-versus-host disease responses. *Blood* **2005**, *106*, 18–26. <https://doi.org/10.1182/blood-2004-08-2996>.
46. López-Pérez, I.C.; Collantes-Fernández, E.; Rojo-Montejo, S.; Navarro-Lozano, V.; Risco-Castillo, V.; Pérez-Pérez, V.; Pereira-Bueno, J.; Ortega-Mora, L.M. Effects of *Neospora caninum* infection at mid-gestation on placenta in a pregnant mouse model. *J. Parasitol.* **2010**, *96*, 1017–1020. <https://doi.org/10.1645/GE-2347.1>.