

**Table S1.** Prevalence of *Toxoplasma gondii* DNA in synanthropic wildlife species of southern Italy according to sex, age, and area, 2020-2022. Missing categories (indicated with “-”) refer to the absence of samples.

Variables	Species		
	Porcupine Pos/Tot (%)	Otter Pos/Tot (%)	Stone marten Pos/Tot (%)
<b>Sex</b>			
Male	1/2 (50.0)	1/3 (33.3)	1/2 (50.0)
Female	1/2 (50.0)	1/2 (50.0)	0/1 (0)
<b>Age</b>			
Juvenile	1/1 (100)	1/2 (50.0)	(-)
Sub-adult	(-)	(-)	(-)
Adult	1/3 (33.3)	1/3 (33.3)	1/3 (33.3)
<b>Area</b>			
Urban	(-)	(-)	1/1 (100)
Peri-urban	1/1 (100)	0/1 (0)	0/2 (0)
Rural	1/3 (33.3)	2/4 (50.0)	(-)

**Table S2.** Prevalence of *Toxoplasma gondii* DNA in synanthropic wildlife species of southern Italy according to the organs examined, 2020-2022. Missing categories (indicated with “-”) refer to the absence of samples.

Species	Organs		
	Heart Pos/Tot (%)	Brain Pos/Tot (%)	Muscle Pos/Tot (%)
<b>Porcupine</b>	1/2 (50.0)	0/2 (0)	1/4 (25.0)
<b>Otter</b>	1/5 (20.0)	1/2 (50.0)	(-)
<b>Beech marten</b>	1/3 (33.3)	1/3 (33.3)	0/3 (0)