




Supplementary Figure 1. Randomization graph—study outline.

Supplementary Table 1. Anthropometric data of patients in the studied groups: the control group, the SE group, and the AoA group.

Parameter	Total n=158 (100%)	Group C n=53 (33.5%)	Group SE n=52 (32.9%)	Group AoA n=53 (33.5%)	p-value
AGE (years)	59±14.1 62 (18)	60.9±13.9 63 (18)	58.7±14.6 61.5 (19.5)	57.4±13.9 62 (16)	p = 0.41
HEIGHT (cm)	167.7±9.4 167 (12)	165.9±9.2 164 (12)	169.5±8.6 170 (12)	167.6±10.1 167 (13)	p = 0.14
WEIGHT (kg)	76.9±16.5 75 (18)	74.5±13.5 75 (15)	79.5±18.8 75.5 (22.5)	76.9±16.9 75 (19)	p = 0.55
BMI	27.3±5.3 27.3 (5.8)	27.1±5 27.4 (4.1)	27.6±6.5 27.2 (5.7)	27.2±4.5 27.8 (6)	p = 0.99
MALE / FEMALE	73 (46.2%) / 85 (53.8%)	23 (43.4%) / 30 (56.6%)	25 (48.1%) / 27 (51.9%)	25 (47.2%) / 28 (52.8%)	p = 0.87

Results are presented as a mean ± SD and median (IQR) for quantitative variables and as numbers (percentages) for nominal variables.

p-values were found by a one-way ANOVA test for quantitative variables; p-values were found by an X2 test for nominal variables.

Abbreviations: Group C: control group; Group SE: Entropy SE group; Group AoA: Adequacy of Anaesthesia group; BMI: body mass index; SD: standard deviation; IQR: interquartile range.