



Article

Lipase Assisted (*S*)-Ketoprofen Resolution from Commercially Available Racemic Mixture

Daniela Estrada-Valenzuela ¹, Víctor H. Ramos-Sánchez ¹, Gerardo Zaragoza-Galán ¹, Jose C. Espinoza-Hicks ¹, Alejandro Bugarin ^{2,*} and David Chávez-Flores ^{1,*}

¹ Facultad de Ciencias Químicas, Universidad Autónoma de Chihuahua, Circuito Universitario, Chihuahua, 31125, Mexico; p291826@uach.mx (D.E.-V.), vramos@uach.mx (V.H.R.-S.), gzaragoza@uach.mx (G.Z.-G.), jhicks@uach.mx (J.C.E.-H.)

² Department of Chemistry & Physics, Florida Gulf Coast University, Fort Myers, FL, Florida, 33965, USA

* Correspondence: abugarin@fgcu.edu (A.B.); dchavezf@uach.mx (D.C.-F.)




Figure S1. (*R,S*)-Ketoprofen FTIR spectrum.




Figure S2. (R,S)-Ketoprofen ¹H NMR spectrum.



Figure S3. Racemic ketoprofen melting point experiment. (m.p. = 93–94 °C).




Figure S4. HPLC chromatogram with the evidence of byproduct formation during the microwave racemization attempt. For this experiment, pure (S)-ketoprofen was used as pure enantiomer to study its racemization using microwave as energy source.




Figure S5. (S)-decyl ketoprofen ester FTIR spectrum (dissolved in methanol).




Figure S6. (S)-decyl ketoprofen ester ^1H NMR spectrum.




Figure S7. (S)-ketoprofen FTIR spectrum.




Figure S8. (S)-ketoprofen ^1H NMR spectrum.