

Article

Tackling Antibiotic Resistance: Influence of Aliphatic Branches on Broad-Spectrum Antibacterial Polytriazoles against ESKAPE Group Pathogens

Cristian Rangel-Núñez ¹, Inmaculada Molina-Pinilla ¹, Cristina Ramírez-Trujillo ¹, Adrián Suárez-Cruz ¹, Samuel Bernal Martínez ² and Manuel Bueno-Martínez ^{1,*}

¹ Departamento de Química Orgánica y Farmacéutica, Facultad de Farmacia, Universidad de Sevilla, C/Profesor García González 2, 41012 Sevilla, Spain

² Servicio de Microbiología, Hospital Universitario Virgen de Valme, 41014 Sevilla, Spain;

* Correspondence: mbueno@us.es




Figure S1. Two-dimensional homonuclear correlation spectrum (COSY) of polymer P2C recorded in deuterochloroform.




Figure S2. Two-dimensional heteronuclear correlation spectrum (HSQC) of polymer P2C recorded in deuteriochloroform.




Figure S3. Two-dimensional homonuclear correlation spectrum (COSY) of polymer P2T recorded in deuteriochloroform.




Figure S4. Two-dimensional heteronuclear correlation spectrum (HSQC) of polymer P2T recorded in deuteriochloroform.