

In Vitro and In Vivo Evaluation of Composite Oral Fast Disintegrating Film: An Innovative Strategy for the Codelivery of Ranitidine HCl and Flurbiprofen

Aisha Rashid ¹, Syed Haroon Khalid ¹, Muhammad Irfan ¹, Sajid Asghar ¹, Waleed Y. Rizg ^{2,3}, Fahad Y. Sabei ⁴, Eman Alfayez ⁵, Hanaa Alkharobi ⁵, Awaji Y. Safhi ⁴, Khaled M. Hosny ³, Muhammad Sohail Arshad ⁶
and Ikram Ullah Khan ^{1,*}

¹ Department of Pharmaceutics, Faculty of Pharmaceutical Sciences, Government College University Faisalabad, Faisalabad 38000, Pakistan; draisharashid@gmail.com (A.R.); haroonkhalid80@gmail.com (S.H.K.); manipharma1@gmail.com (M.I.); sajuhappa@gmail.com (S.A.)

² Center of Innovation in Personalized Medicine (CIPM), 3D Bioprinting Unit, King Abdulaziz University, Jeddah 21589, Saudi Arabia; wrizq@kau.edu.sa

³ Department of Pharmaceutics, Faculty of Pharmacy, King Abdulaziz University, Jeddah 21589, Saudi Arabia; kmhomar@kau.edu.sa

⁴ Department of Pharmaceutics, College of Pharmacy, Jazan University, Jazan 45142, Saudi Arabia; fsabei@jzanu.edu.sa (F.Y.S.); asafhi@jzanu.edu.sa (A.Y.S.)

⁵ Department of Oral Biology, Faculty of Dentistry, King Abdulaziz University, Jeddah 80209, Saudi Arabia; ealfayez@kau.edu.sa (E.A.); halkharobi@kau.edu.sa (H.A.)

⁶ Department of Pharmaceutics, Faculty of Pharmacy, Bahauddin Zakariya University, Multan 60800, Pakistan; sohailarshad@bzu.edu.pk

* Correspondence: ikramglt@gmail.com




Figure S1. Apparent image of (a) M3, (b) L15, (c) F1, and (d) F2.




Figure S2. Optical images showing DT of F1.