

SUPPLEMENTARY MATERIAL

Nanostructured Lipid Carriers Containing Norfloxacin and 2-Aminothiophene Derivative Reduces Fluoroquinolone Resistance in Multidrug-Resistant *Staphylococcus aureus* Strains by Efflux Pump Inhibition

Aléxia Gonçalves Dias ¹, Izabele de Souza Araújo ¹, Rodrigo Santos Aquino de Araújo ², Malu Maria Lucas dos Reis ², Cícera Datiane de Moraes Oliveira Tintino ³, Saulo Relison Tintino ³, Gildênia Alves de Araújo ³, Priscilla Augusta de Sousa Fernandes ¹, Henrique Douglas Melo Coutinho ³, Elquio Eleamen Oliveira ^{2,*} and Francisco Jaime Bezerra Mendonça-Junior ²

¹ Postgraduate Program in Natural and Synthetic Bioactive Products, Federal University of Paraíba, João Pessoa 58051-900, Brazil; alexiajgdias@gmail.com (A.G.D.); izabele.araaujo@gmail.com (I.d.S.A.); prisciasf@gmail.com (P.A.d.S.F.)

² Laboratory of Synthesis and Drug Delivery, Department of Biological Sciences, State University of Paraíba, João Pessoa 58071-160, Brazil; rodrigobiologojp@gmail.com (R.S.A.d.A.); malureisduarte@gmail.com (M.M.L.d.R.); franciscojaime@servidor.uepb.edu.br (F.J.B.M.-J.)

³ Laboratory of Microbiology and Molecular Biology, Universidade Regional do Cariri, Crato 63105-000, Brazil; datianemoraes@gmail.com (C.D.d.M.O.T.); saulo.tintino@urca.br (S.R.T.); gildenia.araujo@urca.br (G.A.d.A.); hdmcoutinho@gmail.com (H.D.M.C.)

* Correspondence: elquioeleamen@servidor.uepb.edu.br; Tel.: +55-84-9987-7523

SUPPLEMENTARY MATERIAL




Figure S1: Calibration curves of peak area versus concentration for norfloxacin (278 nm) and 6CN-Ethyl (221 nm) obtained by HPLC.