SUPPLEMENTARY MATERIAL

Table S1. 2^2 full factorial design of SEDDS mixtures

Comp. / Material	1	2	3	4
Span 20	-1	-1	+1	+1
Span 80	+1	+1	-1	-1
Tween 20	-1	+1	-1	+1
Tween 80	+1	-1	+1	-1

Table S2. Composition of SEDDSs based on the mixture design experimental plan (%, w/w)

1	Miglyol 810 (%)	Tween 80 (%)	Span 80 (%)
1 V	10.00	60.00	30.00
2 V	20.00	60.00	20.00
3 V	10.00	80.00	10.00
4 V	20.00	70.00	10.00
5 C	15.00	67.50	17.50

2	Miglyol 810 (%)	Tween 20 (%)	Span 80 (%)
1 V	10.00	60.00	30.00
2 V	20.00	60.00	20.00
3 V	10.00	80.00	10.00
4 V	20.00	70.00	10.00
5 C	15.00	67.50	17.50

3	Miglyol 810 (%)	Tween 80 (%)	Span 20 (%)
1 V	10.00	60.00	30.00
2 V	20.00	60.00	20.00
3 V	10.00	80.00	10.00
4 V	20.00	70.00	10.00
5 C	15.00	67.50	17.50

4	Miglyol 810 (%)	Tween 20 (%)	Span 20 (%)
1 V	10.00	60.00	30.00
2 V	20.00	60.00	20.00
3 V	10.00	80.00	10.00
4 V	20.00	70.00	10.00
5 C	15.00	67.50	17.50




Figure S1. the applied three-factor-constrained mixture design

Table S3. Droplet size, polydispersity index, zeta potential values and grading of the prepared SEDDS mixtures. ± SD (*n*=3)

Comp.	Droplet size 1 (nm)	Droplet size 2 (nm)	Droplet size 3 (nm)	Mean Droplet size (nm)	Droplet size SD ± (nm)	PDI 1	PDI 2	PDI 3	Mean PDI	PDI SD ±	Zeta potential 1 (mV)	Zeta potential 2 (mV)	Zeta potential 3 (mV)	Mean Zeta potential (mV)	Zeta potential SD ± (mV)	Grade
1																
1 V	365.80	375.10	358.70	366.53	8.225	0.583	0.604	0.569	0.585	0.0176	-15.30	-15.20	-16.10	-15.53	0.493	Grade D
2 V	379.80	380.60	345.50	368.63	20.04	0.458	0.456	0.544	0.486	0.0502	-25.30	-22.30	-24.40	-24.00	1.539	Grade D
3 V	84.48	89.06	82.65	85.40	3.300	0.975	0.990	0.978	0.981	0.0080	-10.20	-9.46	-9.81	-9.82	0.370	Grade B
4 V	34.41	33.26	33.43	33.70	0.621	0.562	0.589	0.565	0.572	0.0150	-12.00	-12.50	-11.90	-12.13	0.321	Grade A
5 C	362.80	362.10	369.60	364.83	4.143	0.440	0.441	0.447	0.443	0.0038	-17.70	-17.90	-18.30	-17.97	0.306	Grade D
2																
1 V	228.00	220.80	226.00	224.93	3.717	0.397	0.457	0.514	0.456	0.0585	-17.80	-17.40	-17.60	-17.60	0.200	Grade D
2 V	95.80	97.05	96.41	96.42	0.625	0.294	0.285	0.299	0.293	0.0071	-14.50	-14.80	-16.00	-15.10	0.794	Grade B
3 V	25.11	24.87	24.96	24.98	0.121	0.293	0.255	0.257	0.268	0.0214	0.26	3.70	3.82	2.59	2.023	Grade A
4 V	100.10	100.50	101.10	100.57	0.503	0.456	0.439	0.465	0.453	0.0132	-9.34	-9.18	-9.62	-9.38	0.223	Grade B
5 C	33.02	33.33	33.42	33.26	0.210	0.391	0.398	0.400	0.396	0.0047	-9.98	-8.85	-7.84	-8.89	1.071	Grade B
3																
1 V	400.60	348.20	367.20	372.00	26.53	0.760	0.802	0.775	0.779	0.0213	-23.20	-34.60	-22.80	-26.87	6.700	Grade D
2 V	475.40	425.10	429.60	443.37	27.83	0.470	0.510	0.511	0.497	0.0234	-23.50	-23.90	-24.00	-23.80	0.265	Grade D
3 V	135.70	127.34	124.56	129.23	3.247	0.961	1.000	0.975	0.979	0.0200	-12.20	-12.00	-11.90	-12.03	0.153	Grade D
4 V	21.82	23.79	22.14	22.58	1.057	0.586	0.550	0.551	0.562	0.0205	9.64	10.20	11.50	10.45	0.954	Grade B
5 C	415.90	412.00	423.10	417.00	5.631	0.490	0.480	0.498	0.489	0.0090	-18.10	-18.50	-18.50	-18.37	0.231	Grade D
4																
1 V	242.10	227.20	241.70	237.00	8.489	0.525	0.597	0.430	0.517	0.0838	-17.90	-17.70	-18.40	-18.00	0.361	Grade D
2 V	43.70	42.55	42.11	42.79	0.821	0.474	0.504	0.503	0.494	0.0170	-14.80	-15.80	-15.70	-15.43	0.551	Grade A
3 V	16.31	16.30	16.33	16.31	0.015	0.222	0.226	0.214	0.221	0.0061	-0.22	-0.03	-0.24	-0.16	0.116	Grade A
4 V	118.70	117.50	113.40	116.53	2.779	0.322	0.314	0.397	0.344	0.0458	-8.75	-9.15	-8.56	-8.82	0.301	Grade B
5 C	19.86	17.97	19.61	19.15	1.027	0.295	0.270	0.299	0.288	0.0157	-3.88	-4.91	-4.36	-4.38	0.515	Grade A




Figure S2. 2D contour plots with the chosen limits of (a) droplet size (blue), (b) PDI (pink) and (c) zeta potential (orange) for the design space in case of composition 1

Table S4. Detailed statistical results of the 3-factor-constrained mixture design for composition 1

Comp. 1 (a) - Factor	Coeffs (recoded comps); Var.:Droplet size (nm); R-sqr=,8909; Adj:,5637 (3 Factor Constrained Mixture in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet size (nm); MS Residual=12732,59					
	Coeff.	**Std. Err.**	**t(1)**	**p**	**-95,% Cnf.Limt**	**+95,% Cnf.Limt**
(A)Miglyol 810	-29.0266	252.2387	-0.115076	0.927061	-3234.02	3175.970
(B)Tween 80	114.2129	108.4804	1.052843	0.483616	-1264.16	1492.587
(C)Span 80	385.1680	111.2894	3.460958	0.179066	-1028.90	1799.234
AC	911.3281	702.7590	1.296786	0.418191	-8018.07	9840.728
Comp. 1 (b) - Factor	**Coeffs (recoded comps); Var.:PDI; R-sqr=,8493; Adj:,3973 (3 Factor Constrained Mixture in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI; MS Residual=,0299919**					
	Coeff.	**Std. Err.**	**t(1)**	**p**	**-95,% Cnf.Limt**	**+95,% Cnf.Limt**
(A)Miglyol 810	0.396200	0.387128	1.023433	0.492628	-4.5227	5.31513
(B)Tween 80	0.971400	0.170804	5.687230	0.110806	-1.1989	3.14167
(C)Span 80	0.537667	0.166493	3.229375	0.191173	-1.5778	2.65315
AB	-0.744000	1.078574	-0.689800	0.615579	-14.4486	12.96058
Comp. 1 (c) - Factor	**Coeffs (recoded comps); Var.:Zeta potetntial (mV); R-sqr=,9862; Adj:,9448 (3 Factor Constrained Mixture in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potetntial (mV); MS Residual=1,671112**					
	Coeff.	**Std. Err.**	**t(1)**	**p**	**-95,% Cnf.Limt**	**+95,% Cnf.Limt**
(A)Miglyol 810	-14.3722	2.889722	-4.9735	0.126317	-51.090	22.34523
(B)Tween 80	-10.1791	1.242784	-8.1906	0.077343	-25.970	5.61193
(C)Span 80	-15.7468	1.274965	-12.3508	0.051433	-31.947	0.45315
AC	-37.4699	8.051018	-4.6541	0.134740	-139.768	64.82798




Figure S3. 2D contour plots with the chosen limits of (a) droplet size (blue), (b) PDI (pink) and (c) zeta potential (orange) for the design space in case of composition 2

Table S5. Detailed statistical results of the 3-factor-constrained mixture design for composition 2

Comp. 2 (a) - Factor	Coeffs (original comps); Var.:Droplet zize (nm); R-sqr=,8703; Adj:,4812 (3 Factor Constrained Mixture in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet zize (nm); MS Residual=3323,411					
	Coeff.	Std. Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	17.51843	13.54200	1.29364	0.418939	-154.549	189.5859
(B)Tween 20	-2.99135	2.18809	-1.36711	0.402049	-30.794	24.8109
(C)Span 80	19.32273	9.44790	2.04519	0.289516	-100.724	139.3697
AC	-1.19991	0.89759	-1.33680	0.408872	-12.605	10.2051
Comp. 2 (b) - Factor	Coeffs (recoded comps); Var.:PDI; R-sqr=,9998; Adj:,9991 (3 Factor Constrained Mixture in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI; MS Residual=,0000073					
	Coeff.	Std. Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	0.129482	0.006030	21.4742	0.029624	0.052868	0.206096
(B)Tween 20	0.267888	0.002660	100.6974	0.006322	0.234085	0.301691
(C)Span 80	0.455258	0.002593	175.5598	0.003626	0.422308	0.488207
AB	1.015030	0.016799	60.4216	0.010535	0.801577	1.228484
Comp. 2 (c) - Factor	Coeffs (recoded comps); Var.:Zeta potetntial (mV); R-sqr=,9994; Adj:,9976 (3 Factor Constrained Mixture in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potetntial (mV); MS Residual=,1466522					
	Coeff.	Std. Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	-21.3737	0.856047	-24.9680	0.025484	-32.2509	-10.4966
(B)Tween 20	2.6981	0.368160	7.3285	0.086336	-1.9799	7.3760
(C)Span 80	-17.5368	0.377694	-46.4312	0.013709	-22.3358	-12.7377
AC	17.9269	2.385021	7.5165	0.084202	-12.3776	48.2315




Figure S4. 2D contour plots with the chosen limits of (a) droplet size (blue), (b) PDI (pink) and (c) zeta potential (orange) for the design space in case of composition 3

Table S6. Detailed statistical results of the 3-factor-constrained mixture design for composition 3

Comp. 3 (a) - Factor	Coeffs (recoded comps); Var.:Droplet size (nm); R-sqr=,8963; Adj:,5854 (3 Factor Constrained Mixture in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet size (nm); MS Residual=14880,33					
	Coeff.	Std. Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	-88.482	272.6840	-0.324484	0.800251	-3553.26	3376.30
(B)Tween 80	160.509	117.2733	1.368671	0.401702	-1329.59	1650.61
(C)Span 20	392.145	120.3100	3.259455	0.189511	-1136.54	1920.83
AC	1327.301	759.7212	1.747090	0.330955	-8325.87	10980.47
Comp 3. (b) - Factor	**Coeffs (recoded comps); Var.:PDI; R-sqr=,8166; Adj:,2665 (3 Factor Constrained Mixture in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI; MS Residual=,0357275**					
	Coeff.	Std. Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	0.225405	0.422527	0.533469	0.688016	-5.1433	5.59412
(B)Tween 80	0.968785	0.186422	5.196736	0.121024	-1.3999	3.33750
(C)Span 20	0.726975	0.181716	4.000601	0.155936	-1.5820	3.03590
AB	-0.388768	1.177197	-0.330249	0.796936	-15.3465	14.56894
Comp. 3 (c) - Factor	**Coeffs (recoded comps); Var.:Zeta potential (mV); R-sqr=,9967; Adj:,9866 (3 Factor Constrained Mixture in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potential (mV); MS Residual=2,948558**					
	Coeff.	Std. Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	33.021	3.83847	8.6027	0.073672	-15.751	81.79358
(B)Tween 80	-12.506	1.65081	-7.5756	0.083552	-33.482	8.46960
(C)Span 20	-27.150	1.69356	-16.0315	0.039659	-48.669	-5.63153
AC	-109.211	10.69431	-10.2120	0.062142	-245.095	26.67361




Figure S5. 2D contour plots with the chosen limits of (a) droplet size (blue), (b) PDI (pink) and (c) zeta potential (orange) for the design space in case of composition 4

Table S7. Detailed statistical results of the 3-factor-constrained mixture design for composition 4

Comp. 4 (a) - Factor	Coeffs (recoded comps); Var.:Droplet size (nm); R-sqr=,9173; Adj:,6694 (3 Factor Constrained Mixture in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet size (nm); MS Residual=2886,633					
	Coeff.	**Std. Err.**	**t(1)**	**p**	**-95,% Cnf.Limt**	**+95,% Cnf.Limt**
(A)Miglyol 810	219.711	120.1017	1.82937	0.318473	-1306.33	1745.748
(B)Tween 20	1.525	51.6522	0.02953	0.981205	-654.78	657.829
(C)Span 20	228.127	52.9897	4.30512	0.145298	-445.17	901.425
AC	-795.512	334.6138	-2.37740	0.253477	-5047.18	3456.159
Comp. 4 (b) - Factor	**Coeffs (recoded comps); Var.:PDI; R-sqr=,8695; Adj:,4781 (3 Factor Constrained Mixture in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI; MS Residual=,0086959**					
	Coeff.	**Std. Err.**	**t(1)**	**p**	**-95,% Cnf.Limt**	**+95,% Cnf.Limt**
(A)Miglyol 810	0.475133	0.208454	2.279322	0.263204	-2.17352	3.123790
(B)Tween 20	0.205267	0.091971	2.231857	0.268168	-0.96334	1.373872
(C)Span 20	0.491667	0.089650	5.484302	0.114819	-0.64744	1.630776
AB	-0.106667	0.580770	-0.183664	0.884365	-7.48605	7.272721
Comp. 4 (c) - Factor	**Coeffs (recoded comps); Var.:Zeta potential (mV); R-sqr=,8766; Adj:,5063 (3 Factor Constrained Mixture in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potential (mV); MS Residual=27,30139**					
	Coeff.	**Std. Err.**	**t(1)**	**p**	**-95,% Cnf.Limt**	**+95,% Cnf.Limt**
(A)Miglyol 810	-17.7653	11.68008	-1.52099	0.370261	-166.175	130.6441
(B)Tween 20	1.2759	5.02326	0.25399	0.841654	-62.551	65.1024
(C)Span 20	-17.1371	5.15333	-3.32544	0.185963	-82.616	48.3422
AC	14.9747	32.54171	0.46017	0.725440	-398.507	428.4563




Figure S6. The acquired design spaces (marked area) for every composition of mixtures and the selected points for validation (1-4, a-d)




Figure S7. The response surface of the effect of HLB value difference on the acquired design space area

Table S8. Detailed statistical results of the 2^2 factorial design to estimate the effect of HLB value difference on the acquired design space area

Factor	Effect Estimates; Var.:Inkscape total%; R-sqr=,9419; Adj:,82571 (Design: 2**(2-0) design (< 200 nm) (Spreadsheet3 in Design space stat) in Design space stat) 2**(2-0) design; MS Residual=44,77617 DV: Inkscape total%									
	Effect	Std. Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt	Coeff.	Std. Err. Coeff.	-95,% Cnf.Limt	+95,% Cnf.Limt
Mean/Interc.	15.2988	3.345750	4.57259	0.137067	-27.213	57.81053	15.2988	3.345750	-27.2130	57.81053
(1)HLB1	14.7255	6.691500	2.20063	0.271531	-70.298	99.74907	7.3628	3.345750	-35.1490	49.87453
(2)HLB2	-22.5635	6.691500	-3.37196	0.183538	-107.587	62.46007	-11.2818	3.345750	-53.7935	31.23003

Table S9. Compositions and predicted values of SEDDS for design space validation

Composition	Code	Component	%	Droplet size (nm)	Droplet size SD ± (nm)	PDI	PDI SD ±	Zeta potential (mV)	Zeta potential (mV) SD ±
C1/Point 1# Replicate 1 and 2	**1D11/1D12**	Miglyol 810	18.5	136.455	89,959	0.557	0,104012	-13.65	5,3338
		Tween 80	71						
		Span 80	10.5						
C1/Point 2# Replicate 1 and 2	**1D21/1D22**	Miglyol 810	19.5	194.109	87,0004	0.4776	0,10059	-15.14	5,1584
		Tween 80	66						
		Span 80	14.5						
C2/Point 1# Replicate 1 and 2	**2D11/2D12**	Miglyol 810	18.5	88.748	166,348	0.3782	0,281737	-11.56	5,0191
		Tween 20	65.5						
		Span 80	16						
C2/Point 2# Replicate 1 and 2	**2D21/2D22**	Miglyol 810	14	134.296	157,954	0.3916	0,267521	-13.35	4,7658
		Tween 20	63.5						
		Span 80	22.5						
C4/Point 1# Replicate 1 and 2	**4D11/4D12**	Miglyol 810	11.5	123.024	250,397	0.3961	0,171323	-10.62	10,3934
		Tween 20	66.5						
		Span 20	22						
C4/Point 2# Replicate 1 and 2	**4D21/4D22**	Miglyol 810	18	72.257	227,035	0.3878	0,155339	-10.27	9,4237
		Tween 20	66						
		Span 20	16						

Table S10. Droplet size, polydispersity index, zeta potential values and grading of the selected validation SEDDS samples. ± SD (*n*=3)

Sample	Droplet size 1 (nm)	Droplet size 2 (nm)	Droplet size 3 (nm)	Mean Droplet size (nm)	Droplet size SD ± (nm)	PDI 1	PDI 2	PDI 3	Mean PDI	PDI SD ±	Zeta potential 1 (mV)	Zeta potential 2 (mV)	Zeta potential 3 (mV)	Mean Zeta potential (mV)	Zeta potential SD ± (mV)	Grade
1D11	161.90	152.80	133.50	149.40	14.502	0.528	0.594	0.647	0.590	0.0596	-7.45	-10.70	-11.80	-9.98	2.262	B
1D12	50.50	50.44	50.17	50.37	0.176	0.936	0.936	0.942	0.938	0.0035	-11.60	-12.60	-12.80	-12.33	0.643	A
1D21	350.70	316.40	322.40	329.83	18.318	0.499	0.510	0.526	0.512	0.0136	-19.60	-20.70	-19.80	-20.03	0.586	C
1D22	349.30	316.20	316.10	327.20	19.139	0.505	0.529	0.524	0.519	0.0127	-19.80	-19.50	-20.40	-19.90	0.458	C
2D11	42.60	41.16	40.93	41.56	0.905	0.311	0.299	0.297	0.302	0.0076	-7.67	-7.62	-8.25	-7.85	0.350	A
2D12	38.29	39.19	39.53	39.00	0.641	0.401	0.372	0.382	0.385	0.0147	-9.02	-9.06	-8.21	-8.76	0.480	A
2D21	468.60	439.60	432.80	447.00	19.013	0.489	0.558	0.548	0.532	0.0373	-0.03	-0.01	0.00	-0.02	0.015	D
2D22	216.10	224.00	227.90	222.67	6.012	0.362	0.387	0.351	0.367	0.0184	-19.90	-14.00	-13.50	-15.80	3.559	C
4D11	101.70	97.11	91.28	96.70	5.222	0.973	0.939	1.000	0.971	0.0306	-0.06	0.05	-0.09	-0.03	0.073	A
4D12	105.10	109.10	106.80	107.00	2.007	0.833	0.861	0.852	0.849	0.0143	-16.60	-18.90	-15.00	-16.83	1.960	A
4D21	30.46	31.47	31.14	31.02	0.515	0.423	0.433	0.425	0.427	0.0053	-11.80	-9.49	-9.19	-10.16	1.428	A
4D22	177.60	231.40	213.50	207.50	27.397	0.282	0.287	0.269	0.279	0.0093	-4.33	-5.46	-5.82	-5.20	0.777	B




Figure S8. Factorial ANOVA comparison of the observed and predicted values (a – droplet size; b – PDI; c – zeta potential) of the validation samples. Data marked with '*' shows significant difference from the predicted value. Vertical bars denote 0.95 confidence intervals.

Table S11. Post Hoc LSD tests of the observed and predicted droplet size, PDI, zeta potential values

LSD test; variable Value (Droplet size dataset in Validation_statistics) Probabilities for Post Hoc Tests Error: Between MS = 12056,, df = 42,000													
Sample	**Dataset**	**{1} 99.885**	**{2} 136.46**	**{3} 328.52**	**{4} 194.11**	**{5} 40.283**	**{6} 88.748**	**{7} 334.83**	**{8} 134.30**	**{9} 101.85**	**{10} 123.02**	**{11} 119.26**	**{12} 72.257**
1D11	**Observed**		0.640056	0.000818	0.231674	0.352486	0.886623	0.000609	0.659881	0.975439	0.767144	0.761370	0.723732
1D11	**Predicted**	0.640056		0.017495	0.523655	0.222331	0.597423	0.014326	0.980898	0.658072	0.881626	0.825810	0.477891
1D21	**Observed**	0.000818	0.017495		0.090759	0.000046	0.003562	0.921100	0.016347	0.000895	0.011391	0.001972	0.001974
1D21	**Predicted**	0.231674	0.523655	0.090759		0.054129	0.246513	0.077053	0.508303	0.241379	0.432282	0.340541	0.181336
2D11	**Observed**	0.352486	0.222331	0.000046	0.054129		0.535851	0.000033	0.232704	0.337018	0.292640	0.219715	0.682561
2D11	**Predicted**	0.886623	0.597423	0.003562	0.246513	0.535851		0.002847	0.614064	0.866815	0.704139	0.696292	0.854939
2D21	**Observed**	0.000609	0.014326	0.921100	0.077053	0.000033	0.002847		0.013369	0.000668	0.009259	0.001486	0.001567
2D21	**Predicted**	0.659881	0.980898	0.016347	0.508303	0.232704	0.614064	0.013369		0.678126	0.900544	0.847387	0.492735
4D11	**Observed**	0.975439	0.658072	0.000895	0.241379	0.337018	0.866815	0.000668	0.678126		0.786382	0.784898	0.705020
4D11	**Predicted**	0.767144	0.881626	0.011391	0.432282	0.292640	0.704139	0.009259	0.900544	0.786382		0.961577	0.574213
4D21	**Observed**	0.761370	0.825810	0.001972	0.340541	0.219715	0.696292	0.001486	0.847387	0.784898	0.961577		0.548153
4D21	**Predicted**	0.723732	0.477891	0.001974	0.181336	0.682561	0.854939	0.001567	0.492735	0.705020	0.574213	0.548153	
LSD test; variable Value (PDI dataset in Validation_statistics) Probabilities for Post Hoc Tests Error: Between MS = ,01840, df = 42,000													
Sample	**Dataset**	**{1} 0.76383**	**{2} 0.55706**	**{3} 0.51550**	**{4} 0.47762**	**{5} 0.34367**	**{6} 0.37820**	**{7} 0.44917**	**{8} 0.39157**	**{9} 0.90967**	**{10} 0.33904**	**{11} 0.35317**	**{12} 0.38780**
1D11	**Observed**		0.036884	0.002837	0.004726	0.000003	0.000236	0.000238	0.000361	0.069600	0.000066	0.000005	0.000321
1D11	**Predicted**	0.036884		0.667023	0.477166	0.031528	0.113823	0.267044	0.142609	0.000666	0.055640	0.039453	0.133960
1D21	**Observed**	0.002837	0.667023		0.694866	0.033816	0.159707	0.401809	0.203414	0.000010	0.072890	0.044369	0.190263
1D21	**Predicted**	0.004726	0.477166	0.694866		0.169911	0.374515	0.768245	0.441588	0.000052	0.217802	0.201566	0.421989
2D11	**Observed**	0.000003	0.031528	0.033816	0.169911		0.720646	0.185180	0.620095	0.000000	0.961750	0.904031	0.647812
2D11	**Predicted**	0.000236	0.113823	0.159707	0.374515	0.720646		0.463492	0.904482	0.000002	0.725440	0.795394	0.931330
2D21	**Observed**	0.000238	0.267044	0.401809	0.768245	0.185180	0.463492		0.551420	0.000001	0.257414	0.227113	0.525795
2D21	**Predicted**	0.000361	0.142609	0.203414	0.441588	0.620095	0.904482	0.551420		0.000003	0.637753	0.690913	0.973008
4D11	**Observed**	0.069600	0.000666	0.000010	0.000052	0.000000	0.000002	0.000001	0.000003		0.000000	0.000000	0.000003
4D11	**Predicted**	0.000066	0.055640	0.072890	0.217802	0.961750	0.725440	0.257414	0.637753	0.000000		0.883611	0.662016
4D21	**Observed**	0.000005	0.039453	0.044369	0.201566	0.904031	0.795394	0.227113	0.690913	0.000000	0.883611		0.719859
4D21	**Predicted**	0.000321	0.133960	0.190263	0.421989	0.647812	0.931330	0.525795	0.973008	0.000003	0.662016	0.719859	

LSD test; variable Value (Zeta potential dataset in Validation_statistics) Probabilities for Post Hoc Tests Error: Between MS = 35,578, df = 42,000													
Sample	**Dataset**	**{1} -11.16**	**{2} -13.65**	**{3} -19.97**	**{4} -15.14**	**{5} -8.305**	**{6} -11.56**	**{7} -7.908**	**{8} -13.35**	**{9} -8.433**	**{10} -10.62**	**{11} -7.682**	**{12} -10.27**
1D11	**Observed**		0.557773	0.014233	0.350598	0.412042	0.923888	0.350657	0.605833	0.433164	0.898998	0.318489	0.834837
1D11	**Predicted**	0.557773		0.141732	0.761292	0.212001	0.670496	0.180615	0.951312	0.222946	0.537110	0.164391	0.491874
1D21	**Observed**	0.014233	0.141732		0.258911	0.001548	0.052865	0.001111	0.124277	0.001720	0.032157	0.000917	0.026596
1D21	**Predicted**	0.350598	0.761292	0.258911		0.112620	0.466881	0.093795	0.715319	0.119311	0.358676	0.084284	0.323434
2D11	**Observed**	0.412042	0.212001	0.001548	0.112620		0.444072	0.908779	0.238230	0.970530	0.586040	0.857235	0.643120
2D11	**Predicted**	0.923888	0.670496	0.052865	0.466881	0.444072		0.391008	0.715426	0.462050	0.847248	0.362614	0.792360
2D21	**Observed**	0.350657	0.180615	0.001111	0.093795	0.908779	0.391008		0.203911	0.879573	0.523763	0.947904	0.577895
2D21	**Predicted**	0.605833	0.951312	0.124277	0.715319	0.238230	0.715426	0.203911		0.250150	0.577864	0.186086	0.530827
4D11	**Observed**	0.433164	0.222946	0.001720	0.119311	0.970530	0.462050	0.879573	0.250150		0.606852	0.828355	0.664807
4D11	**Predicted**	0.898998	0.537110	0.032157	0.358676	0.586040	0.847248	0.523763	0.577864	0.606852		0.489890	0.943652
4D21	**Observed**	0.318489	0.164391	0.000917	0.084284	0.857235	0.362614	0.947904	0.186086	0.828355	0.489890		0.542202
4D21	**Predicted**	0.834837	0.491874	0.026596	0.323434	0.643120	0.792360	0.577895	0.530827	0.664807	0.943652	0.542202	

Table S12. Droplet size, polydispersity index and zeta potential values of the selected HIP loaded SEDDS. ± SD (*n*=3)

Sample	Droplet size 1 (nm)	Droplet size 2 (nm)	Droplet size 3 (nm)	Mean Droplet size (nm)	Droplet size SD ± (nm)	PDI 1	PDI 2	PDI 3	Mean PDI	PDI SD ±	Zeta potential 1 (mV)	Zeta potential 2 (mV)	Zeta potential 3 (mV)	Mean Zeta potential (mV)	Zeta potential SD ± (mV)
1D11															
5 mg/ml	51.25	55.54	50.07	52.29	2.879	0.775	0.714	0.759	0.749	0.0316	-13.2000	-13.5000	-13.1000	-13.2667	0.208
10 mg/ml	49.94	49.29	55.42	51.55	3.367	0.769	0.762	0.723	0.751	0.0248	-11.5000	-11.6000	-11.8000	-11.6333	0.153
15 mg/ml	69.06	56.98	69.68	65.24	7.160	0.650	0.752	0.655	0.686	0.0575	0.0486	-0.0440	-0.0326	-0.0093	0.050
1D21															
5 mg/ml	461.40	460.80	422.30	448.17	22.403	0.545	0.547	0.591	0.561	0.0260	-14.0000	-13.9000	-13.6000	-13.8333	0.208
10 mg/ml	496.00	490.50	488.60	491.70	3.843	0.587	0.578	0.579	0.581	0.0049	-10.6000	-10.6000	-10.9000	-10.7000	0.173
15 mg/ml	315.30	312.20	280.80	302.77	19.087	0.511	0.511	0.609	0.544	0.0566	-0.0101	0.0047	-0.0005	-0.0020	0.008
2D11															
5 mg/ml	83.84	83.97	112.70	93.50	16.625	0.443	0.387	0.264	0.365	0.0916	-0.0593	0.0866	-0.0568	-0.0098	0.084
10 mg/ml	192.50	301.80	350.60	281.63	80.956	0.303	0.362	0.413	0.359	0.0550	0.0044	-0.0526	0.0280	-0.0067	0.041
15 mg/ml	633.20	559.30	686.90	626.47	64.066	0.710	0.634	0.635	0.660	0.0436	-0.0329	-0.0405	-0.0251	-0.0328	0.008
2D22															
5 mg/ml	252.50	240.00	233.40	241.97	9.701	0.386	0.414	0.463	0.421	0.0390	-13.4000	-13.3000	-13.7000	-13.4667	0.208
10 mg/ml	266.80	222.40	257.70	248.97	23.453	0.402	0.478	0.387	0.422	0.0488	0.0103	-0.0671	0.0322	-0.0082	0.052
15 mg/ml	3528.00	3934.00	4303.00	3921.67	387.647	1.000	1.000	1.000	1.000	0.0000	-6.0900	-5.8600	-5.6000	-5.8500	0.245
4D11															
5 mg/ml	136.40	122.20	135.90	131.50	8.058	0.726	0.767	0.723	0.739	0.0246	-14.8000	-15.0000	-15.1000	-14.9667	0.153
10 mg/ml	341.50	351.10	363.90	352.17	11.238	0.555	0.476	0.489	0.507	0.0424	-10.5000	-11.3000	-11.8000	-11.2000	0.656
15 mg/ml	484.10	476.10	391.50	450.57	51.309	0.510	0.507	0.521	0.513	0.0074	-0.0509	0.0823	0.0488	0.0267	0.069
4D21															
5 mg/ml	23.15	23.38	22.75	23.09	0.319	0.322	0.326	0.315	0.321	0.0056	0.0335	-0.0630	-0.0450	-0.0248	0.051
10 mg/ml	34.83	34.53	34.83	34.73	0.173	0.493	0.484	0.491	0.489	0.0047	0.0379	-0.0696	-0.0226	-0.0181	0.054
15 mg/ml	95.43	136.60	232.20	154.74	70.167	0.241	0.275	0.356	0.291	0.0591	-3.2800	-3.4000	-3.4200	-3.3667	0.076

Table S13. Droplet size, PDI and zeta potential values for SEDDS formulations 1D11 and 4D11 in 0.1 M HCl pH 1.4 (1D1G and 4D1G) and in PBS pH 6.8 (1D1P and 4D1P). Indicated values are means of three experiments ± SD (DS-droplet size, PDI-polydispersity index, ZP-zeta potential)

1D1G	**DS**	**PDI**	**ZP**	**4D1G**	**DS**	**PDI**	**ZP**
0 h	49.8±1.7	0.639±0.021	0.79±0.69	**0 h**	78.5±5.4	0.766±0.174	0.23±1.31
1 h	44.7±0.4	0.662±0.009	0.12±0.64	**1 h**	74.2±7.4	0.774±0.173	0.41±0.88
2 h	38.6±1.0	0.649±0.021	0.39±0.32	**2 h**	70.0±0.4	0.832±0.060	0.46±0.73
1D1P	**DS**	**PDI**	**ZP**	**4D1P**	**DS**	**PDI**	**ZP**
0 h	66.9±8.7	0.564±0.078	0.66±2.86	**0 h**	301.3±30.2	0.442±0.008	3.68±0.91
2 h	60.2±6.1	0.542±0.015	3.54±0.62	**2 h**	221.2±34.8	0.460±0.073	4.52±0.88
4 h	59.0±13.2	0.521±0.105	3.42±0.69	**4 h**	156.3±47.0	0.618±0.212	3.92±0.21
6 h	45.7±1.6	0.562±0.024	3.53±0.20	**6 h**	128.2±30.4	0.659±0.200	3.87±0.41
24 h	41.5±0.5	0.526±0.028	3.39±0.41	**24 h**	61.1±3.8	0.849±0.039	4.29±0.44


Detailed statistical results, observes vs. predicted data plots and residual plots of the iterative determination of response surface equations


Composition 1 – Droplet size


Factor	Coeffs (recoded comps); Var.:Droplet size (nm); R-sqr=,7075; Adj:,415 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet size (nm); MS Residual=17072,2					
	Coeff.	Std.Err.	t(2)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	204,5013	204,5163	0,999927	0,422678	-675,461	1084,464
(B)Tween 80	74,3422	120,4632	0,617137	0,600042	-443,969	592,653
(C)Span 80	436,4302	120,4632	3,622935	0,068456	-81,881	954,741

Model	ANOVA; Var.:Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs Sequential fit of models of increasing complexity									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	82589,6	2	41294,82	34144,41	2	17072,20	2,418834	0,292497	0,707503	0,415005
Total Adjusted	116734,1	4	29183,51							

Source	Overall Fit of Model; Var.: Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs				
	SS	df	MS	F	p
Model	82589,6	2	41294,82	2,418834	0,292497
Total Error	34144,4	2	17072,20		
Total Adjusted	116734,1	4	29183,51		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Raw Residuals
150
100
50
0
-50
-100
-150
0
50
100
150
200
250
300
350
400
450
500
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Predicted Values
500
450
400
350
300
250
200
150
100
50
0
-50
0
50
100
150
200
250
300
350
400
450
Observed Values




	Redundant Effects (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) These effects are linear combinations of other effects and cannot be estimated
	Effect
1	BC


Factor	Coeffs (recoded comps); Var.:Droplet size (nm); R-sqr=1, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet size (nm)
	Coeff.
(A)Miglyol 810	-1047,72
(B)Tween 80	83,15
(C)Span 80	366,53
AB	2049,82
AC	2836,92


Model	ANOVA; Var.:Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted

Linear	82589,6	2	41294,82	34144,41	2	17072,20	2,418834	0,292497	0,707503	0,415005
Quadratic	34144,4	2	17072,20	0,00	0	0,00			1,000000	1,000000
Total Adjusted	116734,1	4	29183,51							

Source	Overall Fit of Model; Var.: Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	116734,1	4	29183,51		
Total Error	0,0	0	0,00		
Total Adjusted	116734,1	4	29183,51		







Factor	Coeffs (recoded comps); Var.:Droplet size (nm); R-sqr=,8909; Adj:,5637 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet size (nm); MS Residual=12732,59

	Coeff.	Std.Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	-29,0266	252,2387	-0,115076	0,927061	-3234,02	3175,970
(B)Tween 80	114,2129	108,4804	1,052843	0,483616	-1264,16	1492,587
(C)Span 80	385,1680	111,2894	3,460958	0,179066	-1028,90	1799,234
AC	911,3281	702,7590	1,296786	0,418191	-8018,07	9840,728

Model	ANOVA; Var.:Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	82589,6	2	41294,82	34144,41	2	17072,20	2,418834	0,292497	0,707503	0,415005
Quadratic	21411,8	1	21411,81	12732,59	1	12732,59	1,681654	0,418191	0,890927	0,563706
Total Adjusted	116734,1	4	29183,51							


Source	Overall Fit of Model; Var.: Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	104001,5	3	34667,15	2,722710	0,412729
Total Error	12732,6	1	12732,59		
Total Adjusted	116734,1	4	29183,51		


composition 1 -PDI


Factor	Coeffs (recoded comps); Var.:PDI; R-sqr=,7776; Adj:,5552 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI; MS Residual=,0221314					
	Coeff.	Std.Err.	t(2)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	0,205550	0,232856	0,882735	0,470496	-0,796348	1,207448
(B)Tween 80	0,929550	0,137156	6,777340	0,021085	0,339417	1,519683
(C)Span 80	0,570217	0,137156	4,157444	0,053275	-0,019916	1,160350

Model	ANOVA; Var.:PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs Sequential fit of models of increasing complexity									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	0,154775	2	0,077387	0,044263	2	0,022131	3,496735	0,222384	0,777616	0,555233
Total Adjusted	0,199038	4	0,049759							

Source	Overall Fit of Model; Var.: PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs				
	SS	df	MS	F	p
Model	0,154775	2	0,077387	3,496735	0,222384
Total Error	0,044263	2	0,022131		
Total Adjusted	0,199038	4	0,049759		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Raw Residuals
0,15
0,10
0,05
0,00
-0,05
-0,10
-0,15
-0,20
0,3
0,4
0,5
0,6
0,7
0,8
0,9
1,0
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Predicted Values
1,0
0,9
0,8
0,7
0,6
0,5
0,4
0,3
0,3
0,4
0,5
0,6
0,7
0,8
0,9
1,0
1,1
Observed Values



	Redundant Effects (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) These effects are linear combinations of other effects and cannot be estimated
	Effect
1	BC

Factor	Coeffs (recoded comps); Var.:PDI; R-sqr=1, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI
	Coeff.
(A)Miglyol 810	1,95967
(B)Tween 80	1,00000
(C)Span 80	0,58533
AB	-3,69933
AC	-3,14600

Model	ANOVA; Var.:PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted

Linear	0,154775	2	0,077387	0,044263	2	0,022131	3,496735	0,222384	0,777616	0,555233
Quadratic	0,044263	2	0,022131	0,000000	0	0,000000			1,000000	1,000000
Total Adjusted	0,199038	4	0,049759							


Source	Overall Fit of Model; Var.: PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	0,199038	4	0,049759		
Total Error	0,000000	0	0,000000		
Total Adjusted	0,199038	4	0,049759		


Factor	Coeffs (recoded comps); Var.:PDI; R-sqr=,8493; Adj:,3973 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI; MS Residual=,0299919					
	Coeff.	Std.Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	0,396200	0,387128	1,023433	0,492628	-4,5227	5,31513
(B)Tween 80	0,971400	0,170804	5,687230	0,110806	-1,1989	3,14167
(C)Span 80	0,537667	0,166493	3,229375	0,191173	-1,5778	2,65315
AB	-0,744000	1,078574	-0,689800	0,615579	-14,4486	12,96058


Model	ANOVA; Var.:PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	0,154775	2	0,077387	0,044263	2	0,022131	3,496735	0,222384	0,777616	0,555233
Quadratic	0,014271	1	0,014271	0,029992	1	0,029992	0,475824	0,615579	0,849316	0,397263
Total Adjusted	0,199038	4	0,049759							

Source	Overall Fit of Model; Var.: PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)

	SS	df	MS	F	p
Model	0,169046	3	0,056349	1,878796	0,481538
Total Error	0,029992	1	0,029992		
Total Adjusted	0,199038	4	0,049759		










Composition 1 – Zeta potential

Factor	Coeffs (recoded comps); Var.:Zeta potetntial (mV); R-sqr=,6874; Adj:,3748 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potetntial (mV); MS Residual=18,93391

	Coeff.	Std.Err.	t(2)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	-23,9738	6,810879	-3,51993	0,072092	-53,2787	5,331014
(B)Tween 80	-8,5398	4,011711	-2,12873	0,167059	-25,8008	8,721164
(C)Span 80	-17,8545	4,011711	-4,45060	0,046958	-35,1155	-0,593503


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Raw Residuals
5
4
3
2
1
0
-1
-2
-3
-4
-24
-22
-20
-18
-16
-14
-12
-10
-8
-6
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Predicted Values
-6
-8
-10
-12
-14
-16
-18
-20
-22
-24
-26
-24
-22
-20
-18
-16
-14
-12
-10
-8
Observed Values




	Redundant Effects (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) These effects are linear combinations of other effects and cannot be estimated
	Effect
1	BC


Factor	Coeffs (recoded comps); Var.:Zeta potetntial (mV); R-sqr=1, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potetntial (mV)
	Coeff.
(A)Miglyol 810	-2,7017
(B)Tween 80	-9,8233
(C)Span 80	-15,5333
AB	-23,4833
AC	-59,5300


Model	ANOVA; Var.:Zeta potetntial (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted

Linear	83,2610	2	41,63051	37,86782	2	18,93391	2,19873	0,312624	0,687376	0,374751
Quadratic	36,1967	1	36,19670	1,67111	1	1,67111	21,66025	0,134740	0,986204	0,944815
Total Adjusted	121,1288	4	30,28221							

Source	Overall Fit of Model; Var.: Zeta potetntial (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	119,4577	3	39,81924	23,82799	0,149206
Total Error	1,6711	1	1,67111		
Total Adjusted	121,1288	4	30,28221		









Factor	Coeffs (recoded comps); Var.:Zeta potetntial (mV); R-sqr=,9862; Adj:,9448 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potetntial (mV); MS Residual=1,671112


	Coeff.	Std.Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	-14,3722	2,889722	-4,9735	0,126317	-51,090	22,34523
(B)Tween 80	-10,1791	1,242784	-8,1906	0,077343	-25,970	5,61193
(C)Span 80	-15,7468	1,274965	-12,3508	0,051433	-31,947	0,45315
AC	-37,4699	8,051018	-4,6541	0,134740	-139,768	64,82798

Model	ANOVA; Var.:Zeta potetntial (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs Sequential fit of models of increasing complexity									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	83,2610	2	41,63051	37,86782	2	18,93391	2,198728	0,312624	0,687376	0,374751
Total Adjusted	121,1288	4	30,28221							

Source	Overall Fit of Model; Var.: Zeta potetntial (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs				
	SS	df	MS	F	p
Model	83,2610	2	41,63051	2,198728	0,312624
Total Error	37,8678	2	18,93391		
Total Adjusted	121,1288	4	30,28221		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
0,6
0,4
0,2
0,0
-0,2
-0,4
-0,6
-0,8
-1,0
-1,2
-1,4
-26
-24
-22
-20
-18
-16
-14
-12
-10
-8
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
-8
-10
-12
-14
-16
-18
-20
-22
-24
-26
-26
-24
-22
-20
-18
-16
-14
-12
-10
-8
Observed Values





Composition 2 – Droplet size


Factor	Coeffs (recoded comps); Var.:Droplet zize (nm); R-sqr=,6385; Adj:,2771 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet zize (nm); MS Residual=4631,245					
	Coeff.	Std.Err.	t(2)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	56,3363	106,5202	0,528879	0,649719	-401,983	514,6557
(B)Tween 20	30,1110	62,7420	0,479918	0,678646	-239,846	300,0680
(C)Span 80	188,4150	62,7420	3,003012	0,095301	-81,542	458,3720

Model	ANOVA; Var.:Droplet zize (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs Sequential fit of models of increasing complexity									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	16362,91	2	8181,453	9262,491	2	4631,245	1,766577	0,361457	0,638543	0,277085
Total Adjusted	25625,40	4	6406,349							

Source	Overall Fit of Model; Var.: Droplet zize (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs				
	SS	df	MS	F	p
Model	16362,91	2	8181,453	1,766577	0,361457
Total Error	9262,49	2	4631,245		
Total Adjusted	25625,40	4	6406,349		




Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Predicted Values
Observed Values
0
20
40
60
80
100
120
140
160
180
200
220
240
260



Observed vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Raw Residuals
Observed Values
-80
-60
-40
-20
0
20
40
60
80
0
20
40
60
80
100
120
140
160
180
200
220
240
260


	Redundant Effects (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) These effects are linear combinations of other effects and cannot be estimated
	Effect
1	BC


Factor	Coeffs (recoded comps); Var.:Droplet zize (nm); R-sqr=1, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet zize (nm)
	Coeff.
(A)Miglyol 810	699,78
(B)Tween 20	24,98
(C)Span 80	224,93
AB	-1047,25
AC	-1463,74

Model	ANOVA; Var.:Droplet zize (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	16362,91	2	8181,453	9262,491	2	4631,245	1,766577	0,361457	0,638543	0,277085
Quadratic	9262,49	2	4631,245	0,000	0	0,000			1,000000	1,000000
Total Adjusted	25625,40	4	6406,349							

Source	Overall Fit of Model; Var.: Droplet zize (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	25625,40	4	6406,349		
Total Error	0,00	0	0,000		
Total Adjusted	25625,40	4	6406,349		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
0,010
0,008
0,006
0,004
0,002
0,000
-0,002
-0,004
-0,006
-0,008
-0,010
-0,012
0
20
40
60
80
100
120
140
160
180
200
220
240
260
Predicted Values



Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
260
240
220
200
180
160
140
120
100
80
60
40
20
0
0
20
40
60
80
100
120
140
160
180
200
220
240
260
Observed Values





Factor	Coeffs (recoded comps); Var.:Droplet zize (nm); R-sqr=,8703; Adj:,4812 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet zize (nm); MS Residual=3323,411					
	Coeff.	Std.Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	179,327	128,8681	1,39155	0,396687	-1458,10	1816,751
(B)Tween 20	9,113	55,4224	0,16442	0,896254	-695,10	713,320
(C)Span 80	215,413	56,8575	3,78865	0,164287	-507,03	937,856
AC	-479,963	359,0377	-1,33680	0,408872	-5041,97	4082,043

Model	ANOVA; Var.:Droplet zize (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	16362,91	2	8181,453	9262,491	2	4631,245	1,766577	0,361457	0,638543	0,277085
Quadratic	5939,08	1	5939,080	3323,411	1	3323,411	1,787044	0,408872	0,870308	0,481232
Total Adjusted	25625,40	4	6406,349							

Source	Overall Fit of Model; Var.: Droplet zize (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	22301,99	3	7433,995	2,236857	0,448415
Total Error	3323,41	1	3323,411		
Total Adjusted	25625,40	4	6406,349		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
30
20
10
0
-10
-20
-30
-40
-50
-60
-50
0
50
100
150
200
250
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
250
200
150
100
50
0
-50
0
20
40
60
80
100
120
140
160
180
200
220
240
260
Observed Values




Composition 2 -PDI


Factor	Coeffs (recoded comps); Var.:PDI; R-sqr=,151; Adj:0, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI; MS Residual=,0132847					
	Coeff.	Std.Err.	t(2)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	0,389583	0,180409	2,159444	0,163433	-0,386654	1,165821
(B)Tween 20	0,324983	0,106264	3,058272	0,092346	-0,132232	0,782199
(C)Span 80	0,410850	0,106264	3,866325	0,060854	-0,046366	0,868066


Model	ANOVA; Var.:PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs Sequential fit of models of increasing complexity									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	0,004726	2	0,002363	0,026569	2	0,013285	0,177857	0,849000	0,151000	0,00
Total Adjusted	0,031295	4	0,007824							

Source	Overall Fit of Model; Var.: PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs

	SS	df	MS	F	p
Model	0,004726	2	0,002363	0,177857	0,849000
Total Error	0,026569	2	0,013285		
Total Adjusted	0,031295	4	0,007824		








Redundant Effects (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) These effects are linear combinations of other effects and cannot be estimated


	Effect
1	BC


Factor	Coeffs (recoded comps); Var.:PDI; R-sqr=1, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI
	Coeff.
(A)Miglyol 810	0,153833
(B)Tween 20	0,268333
(C)Span 80	0,456000
AB	0,969000
AC	-0,049000

Model	ANOVA; Var.:PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	0,004726	2	0,002363	0,026569	2	0,013285	0,177857	0,849000	0,151000	0,000000
Quadratic	0,026569	2	0,013285	0,000000	0	0,000000			1,000000	1,000000
Total Adjusted	0,031295	4	0,007824							

Source	Overall Fit of Model; Var.: PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	0,031295	4	0,007824		
Total Error	0,000000	0	0,000000		
Total Adjusted	0,031295	4	0,007824		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
0,010
0,008
0,006
0,004
0,002
0,000
-0,002
-0,004
-0,006
-0,008
-0,010
-0,012
0,24
0,26
0,28
0,30
0,32
0,34
0,36
0,38
0,40
0,42
0,44
0,46
0,48
Predicted Values



Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
0,48
0,46
0,44
0,42
0,40
0,38
0,36
0,34
0,32
0,30
0,28
0,26
0,24
0,24
0,26
0,28
0,30
0,32
0,34
0,36
0,38
0,40
0,42
0,44
0,46
0,48
Observed Values





Factor	Coeffs (recoded comps); Var.:PDI; R-sqr=,9998; Adj:,9991 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI; MS Residual=,0000073					
	Coeff.	Std.Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	0,129482	0,006030	21,4742	0,029624	0,052868	0,206096
(B)Tween 20	0,267888	0,002660	100,6974	0,006322	0,234085	0,301691
(C)Span 80	0,455258	0,002593	175,5598	0,003626	0,422308	0,488207
AB	1,015030	0,016799	60,4216	0,010535	0,801577	1,228484

Model	ANOVA; Var.:PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	0,004726	2	0,002363	0,026569	2	0,013285	0,178	0,849000	0,151000	0,000000
Quadratic	0,026562	1	0,026562	0,000007	1	0,000007	3650,764	0,010535	0,999768	0,999070
Total Adjusted	0,031295	4	0,007824							

Source	Overall Fit of Model; Var.: PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	0,031288	3	0,010429	1433,418	0,019413
Total Error	0,000007	1	0,000007		
Total Adjusted	0,031295	4	0,007824		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
0,0015
0,0010
0,0005
0,0000
-0,0005
-0,0010
-0,0015
-0,0020
-0,0025
-0,0030
0,24
0,26
0,28
0,30
0,32
0,34
0,36
0,38
0,40
0,42
0,44
0,46
0,48
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
0,48
0,46
0,44
0,42
0,40
0,38
0,36
0,34
0,32
0,30
0,28
0,26
0,24
0,24
0,26
0,28
0,30
0,32
0,34
0,36
0,38
0,40
0,42
0,44
0,46
0,48
Observed Values





Composition 2 – Zeta potential


Factor	Coeffs (recoded comps); Var.:Zeta potetntial (mV); R-sqr=,9654; Adj:,9307 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potetntial (mV); MS Residual=4,216057					
	Coeff.	Std.Err.	t(2)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	-16,7800	3,213929	-5,22101	0,034782	-30,6084	-2,95155
(B)Tween 20	1,9138	1,893053	1,01094	0,418461	-6,2314	10,05892
(C)Span 80	-16,5284	1,893053	-8,73106	0,012865	-24,6735	-8,38322

Model	ANOVA; Var.:Zeta potetntial (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs Sequential fit of models of increasing complexity									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	235,0030	2	117,5015	8,432115	2	4,216057	27,87000	0,034638	0,965362	0,930724
Total Adjusted	243,4351	4	60,8588							

Source	Overall Fit of Model; Var.: Zeta potetntial (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs				
	SS	df	MS	F	p
Model	235,0030	2	117,5015	27,87000	0,034638
Total Error	8,4321	2	4,2161		
Total Adjusted	243,4351	4	60,8588		








Redundant Effects (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) These effects are linear combinations of other effects and cannot be estimated


	Effect
1	BC


Factor	Coeffs (recoded comps); Var.:Zeta potetntial (mV); R-sqr=1, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potetntial (mV)
	Coeff.
(A)Miglyol 810	-24,8310
(B)Tween 20	2,5927
(C)Span 80	-17,6000
AB	6,9567
AC	24,4620

Model	ANOVA; Var.:Zeta potetntial (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	235,0030	2	117,5015	8,432115	2	4,216057	27,87000	0,034638	0,965362	0,930724
Quadratic	8,4321	2	4,2161	0,000000	0	0,000000			1,000000	1,000000
Total Adjusted	243,4351	4	60,8588							

Source	Overall Fit of Model; Var.: Zeta potetntial (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	243,4351	4	60,85879		
Total Error	0,0000	0	0,00000		
Total Adjusted	243,4351	4	60,85879		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
0,010
0,008
0,006
0,004
0,002
0,000
-0,002
-0,004
-0,006
-0,008
-0,010
-0,012
-20
-15
-10
-5
0
5
Predicted Values



Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
5
0
-5
-10
-15
-20
-20
-15
-10
-5
0
5
Observed Values





Factor	Coeffs (recoded comps); Var.:Zeta potetntial (mV); R-sqr=,9994; Adj:,9976 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potetntial (mV); MS Residual=,1466522					
	Coeff.	Std.Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	-21,3737	0,856047	-24,9680	0,025484	-32,2509	-10,4966
(B)Tween 20	2,6981	0,368160	7,3285	0,086336	-1,9799	7,3760
(C)Span 80	-17,5368	0,377694	-46,4312	0,013709	-22,3358	-12,7377
AC	17,9269	2,385021	7,5165	0,084202	-12,3776	48,2315

Model	ANOVA; Var.:Zeta potetntial (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	235,0030	2	117,5015	8,432115	2	4,216057	27,87000	0,034638	0,965362	0,930724
Quadratic	8,2855	1	8,2855	0,146652	1	0,146652	56,49738	0,084202	0,999398	0,997590
Total Adjusted	243,4351	4	60,8588							

Source	Overall Fit of Model; Var.: Zeta potetntial (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	243,2885	3	81,09616	552,9831	0,031248
Total Error	0,1467	1	0,14665		
Total Adjusted	243,4351	4	60,85879		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
0,4
0,3
0,2
0,1
0,0
-0,1
-0,2
-20
-15
-10
-5
0
5
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
5
0
-5
-10
-15
-20
-20
-15
-10
-5
0
5
Observed Values





Composition 3 – Droplet size


Factor	Coeffs (recoded comps); Var.:Droplet size (nm); R-sqr=,5799; Adj:,1599 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet size (nm); MS Residual=30149,94					
	Coeff.	Std.Err.	t(2)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	251,6392	271,7855	0,925874	0,452255	-917,759	1421,038
(B)Tween 80	102,4392	160,0858	0,639902	0,587758	-586,354	791,233
(C)Span 20	466,8058	160,0858	2,915974	0,100235	-221,988	1155,599

Model	ANOVA; Var.:Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs Sequential fit of models of increasing complexity									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	83248,9	2	41624,45	60299,88	2	30149,94	1,380581	0,420065	0,579935	0,159869
Total Adjusted	143548,8	4	35887,19							

Source	Overall Fit of Model; Var.: Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs				
	SS	df	MS	F	p
Model	83248,9	2	41624,45	1,380581	0,420065
Total Error	60299,9	2	30149,94		
Total Adjusted	143548,8	4	35887,19		








Redundant Effects (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) These effects are linear combinations of other effects and cannot be estimated
Effect


1	BC


Factor	Coeffs (recoded comps); Var.:Droplet size (nm); R-sqr=1, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet size (nm)
	Coeff.
(A)Miglyol 810	-1189,75
(B)Tween 80	126,93
(C)Span 20	372,00
AB	2215,97
AC	3408,97

Model	ANOVA; Var.:Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	83248,9	2	41624,45	60299,88	2	30149,94	1,380581	0,420065	0,579935	0,159869
Quadratic	60299,9	2	30149,94	0,00	0	0,00			1,000000	1,000000
Total Adjusted	143548,8	4	35887,19							

Source	Overall Fit of Model; Var.: Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	143548,8	4	35887,19		
Total Error	0,0	0	0,00		
Total Adjusted	143548,8	4	35887,19		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
0,010
0,008
0,006
0,004
0,002
0,000
-0,002
-0,004
-0,006
-0,008
-0,010
-0,012
-50
0
50
100
150
200
250
300
350
400
450
500
Predicted Values



Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
500
450
400
350
300
250
200
150
100
50
0
-50
-50
0
50
100
150
200
250
300
350
400
450
500
Observed Values





Factor	Coeffs (recoded comps); Var.:Droplet size (nm); R-sqr=,8963; Adj:,5854 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet size (nm); MS Residual=14880,33					
	Coeff.	Std.Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	-88,482	272,6840	-0,324484	0,800251	-3553,26	3376,30
(B)Tween 80	160,509	117,2733	1,368671	0,401702	-1329,59	1650,61
(C)Span 20	392,145	120,3100	3,259455	0,189511	-1136,54	1920,83
AC	1327,301	759,7212	1,747090	0,330955	-8325,87	10980,47

Model	ANOVA; Var.:Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	83248,9	2	41624,45	60299,88	2	30149,94	1,380581	0,420065	0,579935	0,159869
Quadratic	45419,5	1	45419,55	14880,33	1	14880,33	3,052322	0,330955	0,896340	0,585358
Total Adjusted	143548,8	4	35887,19							

Source	Overall Fit of Model; Var.: Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	128668,4	3	42889,48	2,882294	0,402740
Total Error	14880,3	1	14880,33		
Total Adjusted	143548,8	4	35887,19		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
140
120
100
80
60
40
20
0
-20
-40
-60
-50
0
50
100
150
200
250
300
350
400
450
500
550
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
550
500
450
400
350
300
250
200
150
100
50
0
-50
-50
0
50
100
150
200
250
300
350
400
450
500
Observed Values




Composition 3 -PDI


Factor	Coeffs (recoded comps); Var.:PDI; R-sqr=,7966; Adj:,5933 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI; MS Residual=,019812					
	Coeff.	Std.Err.	t(2)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	0,125783	0,220317	0,570920	0,625652	-0,822163	1,073730
(B)Tween 80	0,946917	0,129770	7,296890	0,018268	0,388562	1,505271
(C)Span 20	0,743983	0,129770	5,733096	0,029103	0,185629	1,302338


Model	ANOVA; Var.:PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs Sequential fit of models of increasing complexity									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	0,155219	2	0,077609	0,039624	2	0,019812	3,917285	0,203364	0,796636	0,593271
Total Adjusted	0,194843	4	0,048711							

Source	Overall Fit of Model; Var.: PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs

	SS	df	MS	F	p
Model	0,155219	2	0,077609	3,917285	0,203364
Total Error	0,039624	2	0,019812		
Total Adjusted	0,194843	4	0,048711		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Raw Residuals
0,10
0,05
0,00
-0,05
-0,10
-0,15
-0,20
-0,25
0,3
0,4
0,5
0,6
0,7
0,8
0,9
1,0
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Predicted Values
1,0
0,9
0,8
0,7
0,6
0,5
0,4
0,3
0,4
0,5
0,6
0,7
0,8
0,9
1,0
1,1
Observed Values




	Redundant Effects (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) These effects are linear combinations of other effects and cannot be estimated
	Effect
1	BC


Factor	Coeffs (recoded comps); Var.:PDI; R-sqr=1, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI
	Coeff.
(A)Miglyol 810	1,93183
(B)Tween 80	1,00000
(C)Span 20	0,77900
AB	-3,61433
AC	-3,43367


Model	ANOVA; Var.:PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted

Linear	0,155219	2	0,077609	0,039624	2	0,019812	3,917285	0,203364	0,796636	0,593271
Quadratic	0,039624	2	0,019812	0,000000	0	0,000000			1,000000	1,000000
Total Adjusted	0,194843	4	0,048711							

Source	Overall Fit of Model; Var.: PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	0,194843	4	0,048711		
Total Error	0,000000	0	0,000000		
Total Adjusted	0,194843	4	0,048711		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
0,010
0,008
0,006
0,004
0,002
0,000
-0,002
-0,004
-0,006
-0,008
-0,010
-0,012
0,4
0,5
0,6
0,7
0,8
0,9
1,0
1,1
Predicted Values



Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
1,1
1,0
0,9
0,8
0,7
0,6
0,5
0,4
0,4
0,5
0,6
0,7
0,8
0,9
1,0
1,1
Observed Values





Factor	Coeffs (recoded comps); Var.:PDI; R-sqr=,8166; Adj:,2665 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI; MS Residual=,0357275					
	Coeff.	Std.Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	0,225405	0,422527	0,533469	0,688016	-5,1433	5,59412
(B)Tween 80	0,968785	0,186422	5,196736	0,121024	-1,3999	3,33750
(C)Span 20	0,726975	0,181716	4,000601	0,155936	-1,5820	3,03590
AB	-0,388768	1,177197	-0,330249	0,796936	-15,3465	14,56894

Model	ANOVA; Var.:PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	0,155219	2	0,077609	0,039624	2	0,019812	3,917285	0,203364	0,796636	0,593271
Quadratic	0,003897	1	0,003897	0,035727	1	0,035727	0,109064	0,796936	0,816634	0,266537
Total Adjusted	0,194843	4	0,048711							

Source	Overall Fit of Model; Var.: PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	0,159115	3	0,053038	1,484528	0,528063
Total Error	0,035727	1	0,035727		
Total Adjusted	0,194843	4	0,048711		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
0,10
0,05
0,00
-0,05
-0,10
-0,15
-0,20
0,4
0,5
0,6
0,7
0,8
0,9
1,0
1,1
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
1,1
1,0
0,9
0,8
0,7
0,6
0,5
0,4
0,4
0,5
0,6
0,7
0,8
0,9
1,0
1,1
Observed Values





Composition 3 – Zeta potential


Factor	Coeffs (recoded comps); Var.:Zeta potential (mV); R-sqr=,6481; Adj:,2961 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potential (mV); MS Residual=155,2199					
	Coeff.	Std.Err.	t(2)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	5,0360	19,50099	0,25824	0,820365	-78,8700	88,94201
(B)Tween 80	-7,7280	11,48638	-0,67280	0,570399	-57,1499	41,69391
(C)Span 20	-33,2933	11,48638	-2,89851	0,101269	-82,7152	16,12857

Model	ANOVA; Var.:Zeta potential (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs Sequential fit of models of increasing complexity									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	571,6495	2	285,8247	310,4398	2	155,2199	1,841418	0,351937	0,648063	0,296126
Total Adjusted	882,0893	4	220,5223							

Source	Overall Fit of Model; Var.: Zeta potential (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs				
	SS	df	MS	F	p
Model	571,6495	2	285,8247	1,841418	0,351937
Total Error	310,4398	2	155,2199		
Total Adjusted	882,0893	4	220,5223		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Raw Residuals
15
10
5
0
-5
-10
-15
-40
-35
-30
-25
-20
-15
-10
-5
0
5
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Predicted Values
5
0
-5
-10
-15
-20
-25
-30
-35
-40
-35
-30
-25
-20
-15
-10
-5
0
5
10
15
Observed Values




	Redundant Effects (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) These effects are linear combinations of other effects and cannot be estimated
	Effect
1	BC


Factor	Coeffs (recoded comps); Var.:Zeta potential (mV); R-sqr=1, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potential (mV)
	Coeff.
(A)Miglyol 810	48,523
(B)Tween 80	-12,033
(C)Span 20	-26,867
AB	-31,193
AC	-138,513


Model	ANOVA; Var.:Zeta potential (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted

Linear	571,6495	2	285,8247	310,4398	2	155,2199	1,841418	0,351937	0,648063	0,296126
Quadratic	310,4398	2	155,2199	0,0000	0	0,0000			1,000000	1,000000
Total Adjusted	882,0893	4	220,5223							

Source	Overall Fit of Model; Var.: Zeta potential (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	882,0893	4	220,5223		
Total Error	0,0000	0	0,0000		
Total Adjusted	882,0893	4	220,5223		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
0,010
0,008
0,006
0,004
0,002
0,000
-0,002
-0,004
-0,006
-0,008
-0,010
-0,012
-35
-30
-25
-20
-15
-10
-5
0
5
10
15
Predicted Values



Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
15
10
5
0
-5
-10
-15
-20
-25
-30
-35
-35
-30
-25
-20
-15
-10
-5
0
5
10
15
Observed Values





Factor	Coeffs (recoded comps); Var.:Zeta potential (mV); R-sqr=,9967; Adj:,9866 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potential (mV); MS Residual=2,948558					
	Coeff.	Std.Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	33,021	3,83847	8,6027	0,073672	-15,751	81,79358
(B)Tween 80	-12,506	1,65081	-7,5756	0,083552	-33,482	8,46960
(C)Span 20	-27,150	1,69356	-16,0315	0,039659	-48,669	-5,63153
AC	-109,211	10,69431	-10,2120	0,062142	-245,095	26,67361

Model	ANOVA; Var.:Zeta potential (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	571,6495	2	285,8247	310,4398	2	155,2199	1,8414	0,351937	0,648063	0,296126
Quadratic	307,4913	1	307,4913	2,9486	1	2,9486	104,2853	0,062142	0,996657	0,986629
Total Adjusted	882,0893	4	220,5223							

Source	Overall Fit of Model; Var.: Zeta potential (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	879,1407	3	293,0469	99,38653	0,073573
Total Error	2,9486	1	2,9486		
Total Adjusted	882,0893	4	220,5223		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
1,0
0,5
0,0
-0,5
-1,0
-1,5
-2,0
-35
-30
-25
-20
-15
-10
-5
0
5
10
15
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
15
10
5
0
-5
-10
-15
-20
-25
-30
-35
-35
-30
-25
-20
-15
-10
-5
0
5
10
15
Observed Values





Composition 4 – Droplet size


Factor	Coeffs (recoded comps); Var.:Droplet size (nm); R-sqr=,4502; Adj:0, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet size (nm); MS Residual=9601,01					
	Coeff.	Std.Err.	t(2)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	15,8610	153,3704	0,103416	0,927068	-644,038	675,7605
(B)Tween 20	36,3290	90,3375	0,402148	0,726482	-352,362	425,0198
(C)Span 20	183,3797	90,3375	2,029940	0,179489	-205,311	572,0704

Model	ANOVA; Var.:Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs Sequential fit of models of increasing complexity									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	15723,62	2	7861,812	19202,02	2	9601,010	0,818853	0,549797	0,450203	0,00
Total Adjusted	34925,64	4	8731,411							

Source	Overall Fit of Model; Var.: Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs				
	SS	df	MS	F	p
Model	15723,62	2	7861,812	0,818853	0,549797
Total Error	19202,02	2	9601,010		
Total Adjusted	34925,64	4	8731,411		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Raw Residuals
120
100
80
60
40
20
0
-20
-40
-60
-80
-100
0
20
40
60
80
100
120
140
160
180
200
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Predicted Values
200
180
160
140
120
100
80
60
40
20
0
-50
0
50
100
150
200
250
300
Observed Values




	Redundant Effects (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) These effects are linear combinations of other effects and cannot be estimated
	Effect
1	BC


Factor	Coeffs (recoded comps); Var.:Droplet size (nm); R-sqr=1, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet size (nm)
	Coeff.
(A)Miglyol 810	704,76
(B)Tween 20	16,31
(C)Span 20	237,00
AB	-976,01
AC	-1712,37


Model	ANOVA; Var.:Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted

Linear	15723,62	2	7861,812	19202,02	2	9601,010	0,818853	0,549797	0,450203	0,000000
Quadratic	19202,02	2	9601,010	0,00	0	0,000			1,000000	1,000000
Total Adjusted	34925,64	4	8731,411							

Source	Overall Fit of Model; Var.: Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	34925,64	4	8731,411		
Total Error	0,00	0	0,000		
Total Adjusted	34925,64	4	8731,411		









Factor	Coeffs (recoded comps); Var.:Droplet size (nm); R-sqr=,9173; Adj:,6694 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Droplet size (nm); MS Residual=2886,633


	Coeff.	Std.Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	219,711	120,1017	1,82937	0,318473	-1306,33	1745,748
(B)Tween 20	1,525	51,6522	0,02953	0,981205	-654,78	657,829
(C)Span 20	228,127	52,9897	4,30512	0,145298	-445,17	901,425
AC	-795,512	334,6138	-2,37740	0,253477	-5047,18	3456,159

Model	ANOVA; Var.:Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	15723,62	2	7861,81	19202,02	2	9601,010	0,818853	0,549797	0,450203	0,000000
Quadratic	16315,39	1	16315,39	2886,63	1	2886,633	5,652046	0,253477	0,917349	0,669397
Total Adjusted	34925,64	4	8731,41							

Source	Overall Fit of Model; Var.: Droplet size (nm) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	32039,01	3	10679,67	3,699697	0,360938
Total Error	2886,63	1	2886,63		
Total Adjusted	34925,64	4	8731,41		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
30
20
10
0
-10
-20
-30
-40
-50
-60
-50
0
50
100
150
200
250
300
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
300
250
200
150
100
50
0
-50
-50
0
50
100
150
200
250
300
Observed Values




Composition 4 – PDI


Factor	Coeffs (recoded comps); Var.:PDI; R-sqr=,8651; Adj:,7302 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI; MS Residual=,0044946					
	Coeff.	Std.Err.	t(2)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	0,447800	0,104937	4,267323	0,050769	-0,003707	0,899307
(B)Tween 20	0,199267	0,061809	3,223886	0,084236	-0,066678	0,465211
(C)Span 20	0,496333	0,061809	8,030054	0,015157	0,230389	0,762278


Model	ANOVA; Var.:PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs Sequential fit of models of increasing complexity									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	0,057655	2	0,028828	0,008989	2	0,004495	6,413850	0,134883	0,865117	0,730235
Total Adjusted	0,066645	4	0,016661							

Source	Overall Fit of Model; Var.: PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs

	SS	df	MS	F	p
Model	0,057655	2	0,028828	6,413850	0,134883
Total Error	0,008989	2	0,004495		
Total Adjusted	0,066645	4	0,016661		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Raw Residuals
0,04
0,02
0,00
-0,02
-0,04
-0,06
-0,08
-0,10
0,15
0,20
0,25
0,30
0,35
0,40
0,45
0,50
0,55
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Linear
Predicted Values
0,55
0,50
0,45
0,40
0,35
0,30
0,25
0,20
0,15
0,15
0,20
0,25
0,30
0,35
0,40
0,45
0,50
0,55
Observed Values




	Redundant Effects (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) These effects are linear combinations of other effects and cannot be estimated
	Effect
1	BC


Factor	Coeffs (recoded comps); Var.:PDI; R-sqr=1, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI
	Coeff.
(A)Miglyol 810	1,31700
(B)Tween 20	0,22067
(C)Span 20	0,51733
AB	-1,69800
AC	-1,69400


Model	ANOVA; Var.:PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted

Linear	0,057655	2	0,028828	0,008989	2	0,004495	6,413850	0,134883	0,865117	0,730235
Quadratic	0,008989	2	0,004495	0,000000	0	0,000000			1,000000	1,000000
Total Adjusted	0,066645	4	0,016661							

Source	Overall Fit of Model; Var.: PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	0,066645	4	0,016661		
Total Error	0,000000	0	0,000000		
Total Adjusted	0,066645	4	0,016661		









Factor	Coeffs (recoded comps); Var.:PDI; R-sqr=,8695; Adj:,4781 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: PDI; MS Residual=,0086959


	Coeff.	Std.Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	0,475133	0,208454	2,279322	0,263204	-2,17352	3,123790
(B)Tween 20	0,205267	0,091971	2,231857	0,268168	-0,96334	1,373872
(C)Span 20	0,491667	0,089650	5,484302	0,114819	-0,64744	1,630776
AB	-0,106667	0,580770	-0,183664	0,884365	-7,48605	7,272721

Model	ANOVA; Var.:PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	0,057655	2	0,028828	0,008989	2	0,004495	6,413850	0,134883	0,865117	0,730235
Quadratic	0,000293	1	0,000293	0,008696	1	0,008696	0,033733	0,884365	0,869519	0,478075
Total Adjusted	0,066645	4	0,016661							

Source	Overall Fit of Model; Var.: PDI (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	0,057949	3	0,019316	2,221312	0,449715
Total Error	0,008696	1	0,008696		
Total Adjusted	0,066645	4	0,016661		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
0,06
0,04
0,02
0,00
-0,02
-0,04
-0,06
-0,08
-0,10
0,15
0,20
0,25
0,30
0,35
0,40
0,45
0,50
0,55
Predicted Values


Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
0,55
0,50
0,45
0,40
0,35
0,30
0,25
0,20
0,15
0,15
0,20
0,25
0,30
0,35
0,40
0,45
0,50
0,55
Observed Values





Composition 4 – Zeta potential


Factor	Coeffs (recoded comps); Var.:Zeta potential (mV); R-sqr=,8504; Adj:,7009 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potential (mV); MS Residual=16,5413					
	Coeff.	Std.Err.	t(2)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	-13,9281	6,366019	-2,18788	0,160172	-41,3188	13,46270
(B)Tween 20	0,6207	3,749681	0,16554	0,883741	-15,5129	16,75429
(C)Span 20	-16,2948	3,749681	-4,34564	0,049087	-32,4284	-0,16121

Model	ANOVA; Var.:Zeta potential (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs Sequential fit of models of increasing complexity									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	188,1089	2	94,05443	33,08261	2	16,54130	5,686035	0,149565	0,850435	0,700869
Total Adjusted	221,1915	4	55,29787							

Source	Overall Fit of Model; Var.: Zeta potential (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs				
	SS	df	MS	F	p
Model	188,1089	2	94,05443	5,686035	0,149565
Total Error	33,0826	2	16,54130		
Total Adjusted	221,1915	4	55,29787		








Redundant Effects (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) These effects are linear combinations of other effects and cannot be estimated
Effect


1	BC


Factor	Coeffs (recoded comps); Var.:Zeta potential (mV); R-sqr=1, (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potential (mV)
	Coeff.
(A)Miglyol 810	-64,9368
(B)Tween 20	-0,1623
(C)Span 20	-18,0000
AB	94,9182
AC	104,1402

Model	ANOVA; Var.:Zeta potential (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	188,1089	2	94,05443	33,08261	2	16,54130	5,686035	0,149565	0,850435	0,700869
Quadratic	33,0826	2	16,54130	0,00000	0	0,00000			1,000000	1,000000
Total Adjusted	221,1915	4	55,29787							

Source	Overall Fit of Model; Var.: Zeta potential (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	221,1915	4	55,29787		
Total Error	0,0000	0	0,00000		
Total Adjusted	221,1915	4	55,29787		


Predicted vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
0,010
0,008
0,006
0,004
0,002
0,000
-0,002
-0,004
-0,006
-0,008
-0,010
-0,012
-20
-18
-16
-14
-12
-10
-8
-6
-4
-2
0
2
Predicted Values



Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
2
0
-2
-4
-6
-8
-10
-12
-14
-16
-18
-20
-20
-18
-16
-14
-12
-10
-8
-6
-4
-2
0
2
Observed Values





Factor	Coeffs (recoded comps); Var.:Zeta potential (mV); R-sqr=,8766; Adj:,5063 (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs DV: Zeta potential (mV); MS Residual=27,30139					
	Coeff.	Std.Err.	t(1)	p	-95,% Cnf.Limt	+95,% Cnf.Limt
(A)Miglyol 810	-17,7653	11,68008	-1,52099	0,370261	-166,175	130,6441
(B)Tween 20	1,2759	5,02326	0,25399	0,841654	-62,551	65,1024
(C)Span 20	-17,1371	5,15333	-3,32544	0,185963	-82,616	48,3422
AC	14,9747	32,54171	0,46017	0,725440	-398,507	428,4563

Model	ANOVA; Var.:Zeta potential (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)									
	SS Effect	df Effect	MS Effect	SS Error	df Error	MS Error	F	p	R-Sqr	R-Sqr Adjusted
Linear	188,1089	2	94,05443	33,08261	2	16,54130	5,686035	0,149565	0,850435	0,700869
Quadratic	5,7812	1	5,78122	27,30139	1	27,30139	0,211755	0,725440	0,876571	0,506285
Total Adjusted	221,1915	4	55,29787							

Source	Overall Fit of Model; Var.: Zeta potential (mV) (3 Factor Constrained Mixture ([No active dataset]) in Experimental_plan_SEDDS) 3 Factor mixture design; Mixture total=100,, 5 Runs (Some terms were excluded from the respective full models)				
	SS	df	MS	F	p
Model	193,8901	3	64,63003	2,367280	0,437940
Total Error	27,3014	1	27,30139		
Total Adjusted	221,1915	4	55,29787		




Observed vs. Predicted Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Predicted Values
Observed Values
4
2
0
-2
-4
-6
-8
-10
-12
-14
-16
-18
-20
-20
-18
-16
-14
-12
-10
-8
-6
-4
-2
0
2


Observed vs. Residual Values
3 Factor mixture design; Mixture total=100,, 5 Runs
Model: Quadratic (some terms were removed from full model)
Raw Residuals
Observed Values
6
5
4
3
2
1
0
-1
-2
-3
-20
-18
-16
-14
-12
-10
-8
-6
-4
-2
0
2