

Supplementary Material

of

Enteral Delivery of Pravastatin Sodium Tablets: Effect of Compounding into a Liquid Form and of Co-Administration of Enteral Nutrition

Serena Logrippo ^{1,2}, Roberta Ganzetti ³, Matteo Sestili⁴, Diego Romano Perinelli⁵, Marco Cespi^{5*} and Giulia Bonacucina⁵.

¹ Hospital Pharmacy, Santa Maria della Stella Hospital, USL Umbria 2, 05018 Orvieto, Italy; serena.logrippo@sanita.marche.it

² Hospital Pharmacy, Engles Profili Hospital, AST Ancona, 60044 Fabriano, Italy;

³ Hospital Pharmacy, Carlo Urbani Hospital, AST Ancona, 60035 Jesi, Italy; roberta.ganzetti@sanita.marche.it

⁴ Territorial Pharmaceutical Service, AST Ancona, 60035 Jesi, Italy; matteo.sestili@sanita.marche.it

⁵ CHIP Building, School of Pharmacy, University of Camerino, via Madonna delle Carceri, 62032 Camerino, Italy; diego.perinelli@unicam.it (D.R.P.); giulia.bonacucina@unicam.it (G.B.)

* Correspondence: marco.cespi@unicam.it; School of Pharmacy, University of Camerino, via Madonna delle Carceri, 62032 Camerino, Italy.

Table S1: Composition of the two nutrition formulae used in the study as provided by the manufacturers.

| Values for 100 mL of product | Nutricomp® Standard Fiber (NSF) | Nutrison® 1Kcal/mol (NWF) |
|------------------------------|---------------------------------|---------------------------|
| Energy (Kcal) | 104 | 100 |
| Protein (g) | 3.8 | 4.0 |
| Fat (g) | 3.3 | 3.9 |
| Saturated | 0.9 | 1 |
| Mono-unsaturated | 1.1 | 2.2 |
| Poly-unsaturated | 1.3 | 0.7 |
| EPA | 0.04 | 0.02 |
| DHA | 0.03 | 0.014 |
| Carbohydrate (g) | 13.8 | 12.3 |
| Sugar | 1.5 | 0.7 |
| Fiber (g) | 1.5 | / |