Supplementary material

Submitted to Special Issue

In vitro evaluation of the antifungal activity of *Trigonella foenum-graecum* seed extract and its potential application in plant protection

Stelica Cristea[1], Alina Perisoara[2*], Bianca-Maria Tihauan[3], Manuela Diana Ene[2], Mariana Constantin[4,5], Alexandru-Mihai Florea[2], Elena Ştefania Ivan[1], Relu Cristinel Zală[1], Bogdan Purcareanu[2,6], Dan Eduard Mihaescu[7] and Lucia Pirvu[8]

[1] University of Agronomical Sciences and Veterinary Medicine, 59 Mărăşti Blvd., District 1, 011464, Bucharest, Romania, stelicacristea@yahoo.com (S.C), elena.ivan@qlab.usamv.ro (E.I.), cristinel.zala@usamv.ro (R.C.Z)

[2] Biotehnos SA., 3-5 Gorunului Street, 075100 Otopeni, Ilfov County, Romania, pavel.alinaa@gmail.com (A.P.), alexandru.florea@biotehnos.com (A.M.F) , bogdanpb89@gmail.com (B.P), diana.ene@biotehnos.com (M.D.E)

[3] Research Institute of the University of Bucharest-ICUB, Splaiul Independenței, no. 95, District 5, Bucharest, Romania, ciubuca.b@gmail.com (B-M.T.)

[4] National Institute for Research & Development in Chemistry and Petrochemistry-ICECHIM, 202 Independentei Spl., 060021 Bucharest, Romania, marriconstantin@yahoo.com

[5] Faculty of Pharmacy, Titu Maiorescu University, 16 Bd. Gh. Șincai, 040441 Bucharest, Romania

[6] Department of Science and Engineering of Oxide Materials and Nanomaterials, National University of Science and Technology Politehnica Bucharest, 011061 Bucharest, Romania, bogdanpb89@gmail.com

[7] Department of Organic Chemistry, National University of Science and Technology Politehnica Bucharest, 011061 Bucharest, Romania, danedmih@gmail.com

[8] National Institute for Chemical-Pharmaceutical Research and Development, District 3, Bucharest, Romania, lucia.pirvu@yahoo.com

* Correspondence: pave.alinaa@gmail.com




Figure S1. Full mass spectra of fenugreek extract using positive ESI ionization




Figure S2. Molecular peak for Protodiosgenina ($C_{51}H_{84}O_{22}$) – m/z calculated is 1049.552, ESI positive




Figure S3. Full mass spectra of Sodium trifluoroacetate using positive ESI ionization for the calibration of the instrument




Figure S4. Full mass spectra of blank sample using positive ESI ionization




Figure S5. Molecular peak for Diosgenina ($C_{27}H_{42}O_3$) – m/z calculated is 415.320, ESI positive




Figure S6. Molecular peak for Trigonelline ($C_7H_7O_2$) – m/z calculated is 138.054, ESI positive




Figure S7. Molecular peak for Tigogenina ($C_{27}H_{44}O_3$) – m/z calculated is 417.336, ESI positive




Figure S8. Molecular peak for Yamogenina ($C_{27}H_{42}O_3$) – m/z calculated is 415.320, ESI positive




Figure S9. Molecular peak for Vitexin ($C_{21}H_{20}O_{10}$) – m/z calculated is 433.112, ESI positive




Figure S10. Molecular peak for Acid salicylic ($C_7H_6O_3$) – m/z calculated is 139.038, ESI positive




Figure S11. Molecular peak for Apigenin 6-C-galactoside 8-C – arabinoside ($C_{26}H_{28}O_{14}$) – m/z calculated is 565.115, ESI positive




Figure S12. Molecular peak for Luteoline – 7 – O glucoside ($C_{21}H_{18}O_{12}$) – m/z calculated is 463.087, ESI positive




Figure S13. Molecular peak for Apigenin – 7 – O – glucoside ($C_{21}H_{20}O_{10}$) – m/z calculated is 433.112, ESI positive




Figure S14. Molecular peak for Acid abscisic ($C_{15}H_{20}O_4$) – m/z calculated is 265.143, ESI positive




Figure S15. Molecular peak for Rutin ($C_{27}H_{30}O_{16}$) – m/z calculated is 611.160, ESI positive




Figure S16. Molecular peak for t-Resveratrol ($C_{14}H_{12}O_3$) – m/z calculated is 228.078, ESI positive