Comparison of Satellite Driven Surface Energy Balance Models in Estimating Crop Evapotranspiration in Semi-Arid to Arid Inter-Mountain Region

Bibek Acharya [1] and Vivek Sharma [1,*]

[1] Agricultural and Biological Engineering Department, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, 32611, USA

* Correspondence: Vivek Sharma, vsharma1@ufl.edu; Tel.: +1-352-294-6725

Supplementary Material:




Figure S1. spatio-temporal distribution of METRIC-estimated ET_{24} (a) 15 May 2018, (b) 08 June 2018, (c) 02 July 2018, (d) 18 July 2018, (e) 26 July 2018, (f) 11 August 2018, (g) 04 September 2018, and (h) September 12, 2018.




Figure S2. spatio-temporal distribution of SEBAL-estimated ET_{24} (a) 15 May 2018, (b) 08 June 2018, (c) 02 July 2018, (d) 18 July 2018, (e) 26 July 2018, (f) 11 August 2018, (g) 04 September 2018, and (h) September 12, 2018.




Figure S3. spatio-temporal distribution of SEBS-estimated ET_{24} (a) 15 May 2018, (b) 08 June 2018, (c) 02 July 2018, (d) 18 July 2018, (e) 26 July 2018, (f) 11 August 2018, (g) 04 September 2018, and (h) September 12, 2018.




Figure S4. spatio-temporal distribution of S-SEBI-estimated ET_{24} (a) 15 May 2018, (b) 08 June 2018, (c) 02 July 2018, (d) 18 July 2018, (e) 26 July 2018, (f) 11 August 2018, (g) 04 September 2018, and (h) September 12, 2018.