Recovery of Bioactive Extracts from Olive Leaves Using Conventional and Microwave-Assisted Extraction with Classical and Deep Eutectic Solvents

Eleni Boli [1,*], Nikos Prinos [1], Vasiliki Louli [1], Georgia Pappa [1], Haralambos Stamatis [2], Kostis Magoulas [1] and Epaminondas Voutsas [1]

[1] Laboratory of Thermodynamics and Transport Phenomena, School of Chemical Engineering, National Technical University of Athens (NTUA), Zografou Campus, 15780 Athens, Greece
[2] Department of Biological Applications and Technologies, Biotechnology Laboratory, University of Ioannina, 45110 Ioannina, Greece
* Correspondence: bolieleni@chemeng.ntua.gr

Supplementary data




Figure S1. HPLC chromatogram of tyrosol (7.8 min), hydroxytyrosol (11.02 min) and oleuropein (24. 78 min) standards.




Figure S2. HPLC chromatogram of OL extract (solvent: ChCl: urea (1:2), T = 55 °C, mass ratio 1:30 w/w, t = 24 h).




Figure S3. HPLC chromatogram of OL extract (solvent: ChCl:AA (1:2), T = 55 °C, mass ratio 1:30 w/w).




Figure S4. HPLC chromatogram of OL extract (solvent: Ethanol = 55 °C, mass ratio 1:20 w/w).