

Table S1. QLQ-C30 by work status.

QLQ-C30	Mean ± SD		p-value ^a
	Non-working	Working	
Functional scale			
Physical functioning (PF)	80.1 ± 22.7	87.7 ± 15.9	*.006
Role functioning (RF)	75.7 ± 28.4	89.3 ± 18.2	**<.001
Emotional functioning (EF)	74.3 ± 27.6	85.0 ± 20.4	*.002
Cognitive functioning (CF)	73.3 ± 26.7	84.3 ± 21.1	**<.001
Social functioning (SF)	65.9 ± 30.8	82.6 ± 24.5	**<.001
Symptom scale			
Fatigue (FA)	32.8 ± 26.8	24.5 ± 23.4	*.013
Nausea/vomiting (NV)	6.4 ± 14.7	4.1 ± 13.8	.055
Pain (PA)	19.1 ± 25.3	13.6 ± 21.2	.054
Dyspnea (DY)	15.7 ± 21.9	8.1 ± 17.2	**<.001
Insomnia (IN)	24.0 ± 33.5	16.6 ± 25.3	.150
Appetite loss (AP)	12.3 ± 24.4	7.2 ± 18.8	.051
Constipation (CO)	17.6 ± 27.9	13.0 ± 23.5	.198
Diarrhea (DI)	12.7 ± 21.6	8.3 ± 20.5	*.022
Financial difficulties (FI)	39.2 ± 37.7	22.1 ± 31.6	**<.001