sustainability

MDPI

Supplementary Material for the Article:

Plant Growth Regulators and Short-Term Irrigation for Berry Maturation Homogeneity and Increased *Coffea arabica* Bean Quality

Miroslava Rakočević [1,*], Eunice Reis Batista [1], Fabio Takeshi Matsunaga [2] and Maria Brígida dos Santos Scholz [3]

[1] Embrapa Meio Ambiente, Jaguariúna 13820–000, SP, Brazil; eunice.reis@embrapa.br
[2] UniSENAI PR – Campus Londrina, Londrina 86062-030, PR, Brazil; ftakematsu@gmail.com
[3] Instituto Agronômico do Paraná, Londrina 86047–902, PR, Brazil; mbrischolz@hotmail.com
* mima.rakocevic61@gmail.com; +55–19–97161–8918




Figure S1. Daily maximum (Tair_max) and minimum (Tair_min) air temperatures and rainfall (Rain) during the period from Arabica coffee flowering to harvest and indication of the dry period (tagged with black oval) when the irrigation was conducted.

Academic Editor: Firstname Last-name

Received: date
Revised: date
Accepted: date
Published: date

Citation: To be added by editorial staff during production.

Copyright: © 2025 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).




Figure S2. Fresh mass (FM) of (**A**) red, (**B**) green, and (**C**) total berries, and (**D**) % of red berries in the total FM after the external application of plant growth regulators (PGRs) (gibberellic acid—GA3, ethylene—Ethephon or water—IRR) over the plant strata (S1, < 80 cm; S2, 81–160 cm; S3, > 160 cm). Mean values ± SE and *p*-values (marked in bold when significant) are shown (n = 4). Inside each figure, different lowercase letters indicate significant difference among PGRs for each stratum, while different uppercase letters indicate significant differences among strata for each PGR.




Figure S3. Fresh mass (FM) of (**A**) red, (**B**) green, and (**C**) total berries and (**D**) % of red berries in the total FM dependent on water regime (irrigated—IRR or rainfed—NI) over plant strata (S1, < 80 cm; S2, 81–160 cm; S3, > 160 cm). Mean values ± SE and *p*-values (marked in bold when significant) are shown (n = 4). Inside each figure, different lowercase letters indicate significant differences between water regimes for each stratum, while different uppercase letters indicate significant differences among strata for each water regime.