sustainable chemistry

MDPI

*Article*

# Accelerated Synthesis of 4-Arylideneisoxazolones in Glycerol Medium

**Jamal Mohammadi Masiri, Hamzeh Kiyani * and Jalal Albadi**

Department of Chemistry, Faculty of Science, Shahrekord University, Shahrekord 8818634141, Iran; jamalmasirii@gmail.com (J.M.M.); albadi@sku.ac.ir (J.A.)
* Correspondence: hkiyani@sku.ac.ir or hakiyani@gmail.com





Figure. S1. $^{1}$H NMR (DMSO-$d_6$, 400 MHz) spectrum of 4-(4-hydroxy-3-methoxybenzylidene)-3-methylisoxazol-5(4*H*)-one (**4a**).





Figure. S2. $^{13}C$ NMR (DMSO-$d_6$, 100 MHz) spectrum of **4a.**

https://doi.org/10.3390/suschem7030039





Figure. S3. $^{1}$H NMR (DMSO-$d_6$, 400 MHz) spectrum of 4-(4-Hydroxybenzylidene)-3-phenylisoxazol-5(4*H*)-one (**4m**).





Figure. S4. $^{13}C$ NMR (DMSO-$d_6$, 100 MHz) spectrum of **4m.**

https://doi.org/10.3390/suschem7030039





Figure. S5. $^{1}$H NMR (DMSO-$d_6$, 400 MHz) spectrum of 4-(3-Ethoxy-4-hydroxybenzylidene)-3-phenylisoxazol-5(4*H*)-one (**4o**).





Figure. S6. The $^{13}C$ NMR (DMSO-$d_6$, 100 MHz) spectrum of **4o**.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.