

Table S4. Detection of atypical CPB2 on field samples with both PCR and sELISA. OD<sub>450</sub> values and PCR outcomes are shown. ND: not determined.

Sample number	Faeces		Intestinal content		Rectal swab	
	sELISA	PCR	sELISA	PCR	sELISA	PCR
1	0.084	Negative	0.087	Negative	0.071	Negative
2	0.067	Negative	0.093	Negative	0.06	Negative
3	0.086	Negative	0.07	Negative	0.063	Negative
4	0.058	Negative	0.053	Negative	0.053	Negative
5	ND	ND	0.061	Negative	ND	ND
6	0.063	Negative	0.077	Negative	0.067	Negative
7	0.057	Negative	0.056	Negative	0.05	Negative
8	0.05	Negative	0.051	Negative	0.048	Negative
9	0.05	Negative	0.05	Negative	0.05	Negative
10	0.049	Negative	0.052	Negative	0.051	Negative
11	0.079	Negative	0.069	Negative	0.064	Negative
12	0.076	Negative	0.065	Negative	0.06	Negative
13	0.08	Negative	0.107	Negative	0.071	Negative
14	ND	ND	0.075	Negative	0.061	Negative
15	0.089	Negative	0.054	Negative	0.08	Negative
16	0.05	Negative	0.071	Negative	0.056	Negative
17	ND	ND	0.093	Negative	0.094	Negative
18	0.054	Negative	0.053	Negative	0.049	Negative
19	0.071	Negative	0.073	Negative	0.086	Negative
20	0.053	Negative	0.076	Negative	0.05	Negative
21	0.053	Negative	0.05	Negative	0.058	Negative
22	0.051	Negative	0.075	Negative	0.051	Negative
23	0.063	Negative	0.069	Negative	0.054	Negative
24	ND	ND	0.078	Negative	ND	ND
25	0.08	Negative	0.055	Negative	0.131	Negative
26	0.306	Negative	0.062	Negative	0.059	Negative
27	0.28	Negative	0.05	Negative	0.124	Negative
28	ND	Negative	0.126	Negative	0.051	Negative
29	0.39	Negative	0.091	Negative	0.209	Negative
30	0.086	Negative	0.183	Negative	0.164	Negative