

Supplementary Materials: COVID-19 Severity and Mortality in Solid Organ Transplantation: Differences between Liver, Heart, and Kidney Recipients

Ricardo Wesley Alberca *, Gabriela Gama Freire Alberca, Lucas Chaves Netto, Raquel Leão Orfali, Sarah Cristina Gozzi-Silva, Alberto José da Silva Duarte, Valeria Aoki, Maria Notomi Sato and Gil Benard

Table S1. Solid organ transplant treatment.

Pre and during COVID-19 treatment	CONTROL	LIVER	KIDNEY	HEART
Tacrolimus	-	6	-	-
Cyclosporine	-	-	1	-
Tacrolimus + mycophenolate mofetil	-	1	5	-
Tacrolimus + azathioprine	-	-	1	-
Tacrolimus + sirolimus	-	-	2	-
Tacrolimus + everolimus	-	-	1	-
Tacrolimus + cyclosporine	-	-	-	1
Mycophenolate mofetil + prednisone	-	-	1	-
Mycophenolate mofetil + cyclosporine	-	-	1	-
Tacrolimus + mycophenolate mofetil + prednisone	-	-	9	1
Tacrolimus + sirolimus + prednisone	-	-	-	1
Mycophenolate mofetil + prednisone + cyclosporine	-	-	4	4
COVID-19 treatment	CONTROL	LIVER	KIDNEY	HEART
Antibiotics	25	7	25	7
Anticoagulants	25	7	25	7
Antivirals	10	1	8	2
Systemic corticosteroids	20	7	22	7