



Figure S1. Plate images of inhibition of infection. Cell monolayer survival was used as a proxy measure for infection. Following pre-treatment with product A, B, or C, and SARS-CoV-2 infection of Vero A/T cells in 24-well plates, cells were fixed with methanol and stained with crystal violet. The plate images show the degree of monolayer staining, under each of the conditions tested, from one representative experiment.

Table S1: Statistical analysis for Figure 1

| Test | MOI | Product | Timepoint | P value | Significance |
|-----------------|------|---------|-----------|---------|--------------|
| No-product vs A | 0.01 | 1g | 24h | <0.0001 | **** |
| | | | 48h | <0.0001 | **** |
| | | | 72h | 0.0052 | ** |
| | | 3g | 24h | 0.0003 | *** |
| | | | 48h | <0.0001 | **** |
| | | | 72h | <0.0001 | **** |
| | | 6.4g | 24h | <0.0001 | **** |
| | | | 48h | 0.0004 | *** |
| | | | 72h | 0.0005 | *** |
| | 1 | 1g | 24h | <0.0001 | **** |
| | | | 48h | 0.0007 | *** |
| | | | 72h | 0.0074 | ** |
| | | 3g | 24h | <0.0001 | **** |
| | | | 48h | <0.0001 | **** |
| | | | 72h | <0.0001 | **** |
| | | 6.4g | 24h | 0.0002 | *** |
| | | | 48h | <0.0001 | **** |
| | | | 72h | <0.0001 | **** |
| No-product vs B | 0.01 | 1g | 24h | <0.0001 | **** |
| | | | 48h | 0.0012 | ** |
| | | | 72h | 0.0011 | ** |
| | | 3g | 24h | <0.0001 | **** |
| | | | 48h | <0.0001 | **** |
| | | | 72h | <0.0001 | **** |
| | | 6.4g | 24h | <0.0001 | **** |
| | | | 48h | 0.0004 | *** |
| | | | 72h | 0.0005 | *** |
| | 1 | 1g | 24h | <0.0001 | **** |
| | | | 48h | 0.0005 | *** |
| | | | 72h | <0.0001 | **** |
| | | 3g | 24h | <0.0001 | **** |
| | | | 48h | 0.0002 | *** |
| | | | 72h | <0.0001 | **** |
| | | 6.4g | 24h | <0.0001 | **** |
| | | | 48h | <0.0001 | **** |
| | | | 72h | <0.0001 | **** |
| No-product vs C | 0.01 | 1g | 24h | <0.0001 | **** |
| | | | 48h | 0.0022 | ** |
| | | | 72h | 0.0069 | ** |
| | | 3g | 24h | <0.0001 | **** |
| | | | 48h | <0.0001 | **** |
| | | | 72h | <0.0001 | **** |
| | | 6.4g | 24h | <0.0001 | **** |
| | | | 48h | 0.0004 | *** |
| | | | 72h | 0.0005 | *** |
| | 1 | 1g | 24h | <0.0001 | **** |
| | | | 48h | 0.0033 | ** |
| | | | 72h | 0.0052 | ** |
| | | 3g | 24h | <0.0001 | **** |
| | | | 48h | 0.0101 | * |
| | | | 72h | 0.0173 | * |
| | | 6.4g | 24h | <0.0001 | **** |
| | | | 48h | <0.0001 | **** |
| | | | 72h | <0.0001 | **** |

Table S2: Statistical analysis for Figure 2

| Virus Strain | Product | Timepoint | <i>P</i> value | Significance |
|---------------------|----------------|------------------|-----------------------|---------------------|
| England-2 | A | 24h | 0.4119 | ns |
| | | 48h | <0.0001 | **** |
| | | 72h | <0.0001 | **** |
| | B | 24h | 0.0415 | * |
| | | 48h | <0.0001 | **** |
| | | 72h | <0.0001 | **** |
| | C | 24h | 0.1145 | ns |
| | | 48h | <0.0001 | **** |
| | | 72h | <0.0001 | **** |
| Alpha | A | 24h | 0.0804 | ns |
| | | 48h | <0.0001 | **** |
| | | 72h | <0.0001 | **** |
| | B | 24h | 0.0368 | * |
| | | 48h | <0.0001 | **** |
| | | 72h | <0.0001 | **** |
| | C | 24h | 0.2315 | ns |
| | | 48h | <0.0001 | **** |
| | | 72h | <0.0001 | **** |
| Beta | A | 24h | 0.0830 | ns |
| | | 48h | 0.0001 | *** |
| | | 72h | <0.0001 | **** |
| | B | 24h | 0.0600 | ns |
| | | 48h | <0.0001 | **** |
| | | 72h | <0.0001 | **** |
| | C | 24h | 0.0944 | ns |
| | | 48h | 0.0002 | *** |
| | | 72h | <0.0001 | **** |