SUPPLEMENTARY INFORMATION

Sequences encoding a novel toursvirus identified from southern and northern corn rootworms (Coleoptera: Chrysomelidae)

Sijun Liu [1], Thomas W. Sappington [2], Brad S. Coates [2], Bryony C. Bonning [3]

Maps of DiTV3a genomic fragments isolated from SCR (Figure S1) and NCR (Figure S2). For both figures, Arrows indicate ORFs and ORF orientation. ORFs that hit toursvirus genes (DjTV2a/DpTV1a) are indicated.

Figure S1. DiTV3a genomic fragments (28) isolated from the southern corn rootworm.



Figure S2. DiTV3a genomic fragments (42) isolated from the northern corn rootworm




Figure S2 ctd.



F22 (1875bp)




F23 (2753bp)




F24 (2211bp)



F25 (1787bp)




F26 (3641bp)




F27 (2750bp)



F28 (2526bp)



F29 (1914bp)



F30 (1425bp)



F31 (1013bp)




F32 (2455bp)



F33 (1274bp)



F34 (673bp)



F35 (1970bp)



F36 (1399bp)




F37 (1401bp)



F38 (1424bp)



F39 (1327bp)



F40 (1305bp)



F41 (1246bp)



F42 (1135bp)