

Article

A BioID-Derived Proximity Interactome for SARS-CoV-2 Proteins

Danielle G. May ¹, Laura Martin-Sancho ^{2†}, Valesca Anschau ², Sophie Liu ³, Rachel J. Chrisopoulos ¹, Kelsey L. Scott ¹, Charles T. Halfmann ¹, Ramon Díaz Peña ², Dexter Pratt ³, Alexandre R. Campos ^{2,*} and Kyle J. Roux ^{1,4,*}

Supplementary Materials




Figure S1. Epifluorescent images of all BioID fusion-protein localizations in A549 cells. A549 human lung cells stably expressing SARS-CoV-2 BioID2 fusion-proteins were assessed for fusion-protein expression and localization (red) and promiscuous biotinylation (green) following the addition of exogenous biotin.




Figure S2. Western blot analysis of A549 cell stably expressing SARS-CoV-2 BioID fusion proteins. A549 cells stably expressing BioID2 fusion-proteins were evaluated for fusion-protein expression (anti-BioID2) and promiscuous biotinylation (streptavidin-HRP) and A549 cells stably expressing TurboID-KDEL control or ORF8-TurboID were evaluated for fusion-protein expression (anti-HA) and promiscuous biotinylation (streptavidin-HRP).