

Canine Distemper Virus Alters Defense Responses in an *Ex Vivo* Model of Pulmonary Infection

Elisa Chludzinski ^{1,3}, Małgorzata Ciurkiewicz ¹, Melanie Stoff ¹, Johanna Klemens ¹, Johannes Krüger ¹, Dai-Lun Shin², Georg Herrler ^{2,*}, Andreas Beineke ^{1,3,*,**}

¹ Department of Pathology, University of Veterinary Medicine Hannover, 30559 Hannover, Germany

² Institute of Virology, University of Veterinary Medicine Hannover, 30559 Hannover, Germany

³ Center for Systems Neuroscience (ZSN), 30559 Hannover, Germany

* Both authors have contributed equally last (G.H., A.B.)

** Correspondence: andreas.beineke@tiho-hannover.de (A.B.)

Supplementary Material:


This file includes: Supplementary Table S1, Supplementary Figure S1

Supplementary Table S1: Details of dogs used for precision-cut lung slices (PCLSs)

Dog No	Age (years)	Sex	Breed
1	13	m	Petit Basset Griffon Vendeen
2	6.5	mc	Labrador
3	18	m	mixed breed
4	8	f	mixed breed
5	12	fs	mixed breed
6	11	fs	Australian Shepherd

m = male; mc = male castrated; f = female; fs = female spayed

Supplementary Figure S1:



Supplementary Figure S1: Scheme for the evaluation of ciliary beating activity by light microscopy. To evaluate the percentage of ciliary beating activity, the bronchus is divided into ten equally-sized segments.