

Supporting Information

Clinical and experimental determination of protection afforded by BCG vaccination against infection with non-tuberculous mycobacteria: a role in Cystic Fibrosis?


Warner *et. al.*

Supplementary Table S1. Monoclonal antibodies used for CD4⁺ T cell analysis.

Marker	Clone	Fluorochrome	Dilution	Manufacturer
B220	PA3-6B2	PerCP/Cy5.5	1 in 200	BD Pharmingen
CD4	RM4-5	AF700	1 in 200	BD Pharmingen
CD44	IM7	PE/Cy7	1 in 300	BD Pharmingen
CD62L	MEL-14	PE	1 in 300	BioLegend
CD8a	53-6.7	APC-Cy7	1 in 200	BD Pharmingen
FcRIII/II	2.4G2	NA	1 in 200	BD Pharmingen
KLRG-1	2F1/KLRG1	FITC	1 in 300	BioLegend
Ror γ T	Q31-378	PECF594	1 in 200	BD Horizon
Tbet	4B10	APC	1 in 200	BioLegend

Supplementary Table S2. Monoclonal antibodies used to assess antigen-specific cytokine production.

Marker	Clone	Fluorochrome	Dilution	Manufacturer
CD4	RM4-5	AF700	1 in 200	BD Pharmingen
CD44	IM7	FITC	1 in 300	BD Pharmingen
CD8a	53-6.7	APC-Cy7	1 in 200	BD Pharmingen
FcRIII/II	2.4G2	NA	1 in 200	BD Pharmingen
IFN γ	XMG1.2	PE/Cy7	1 in 300	BD Pharmingen
IL-17	TC11-18H10	Pacific Blue	1 in 200	BD Horizon
IL-2	JES6-5H4	PE	1 in 200	Miltenyi Biotec
TNF	MP6-XT22	PerCP/Cy5.5	1 in 200	BD Pharmingen



Supplementary Figure S1. Gating strategy for expression of cytokines and transcription factors by CD4⁺ T cells