

Supplementary Table S1. Minimal Inhibitory Concentrations (MIC) in different bacterial isolates and antimicrobials.

Gram-negative												Gram-positive				
Acinetobacter spp			Pseudomonas spp		Bordetella spp		Escherichia spp		Klebsiella spp		Enterobacter spp		Staphylococcus spp		Enterococcus spp	
Ab	MIC ₅₀	MIC ₉₀	MIC ₅₀	MIC ₉₀	MIC ₅₀	MIC ₉₀	MIC ₅₀	MIC ₉₀	MIC ₅₀	MIC ₉₀	MIC ₅₀	MIC ₉₀	MIC ₅₀	MIC ₉₀	MIC ₅₀	MIC ₉₀
AMP	16	16	16	32	16	16	16	32	16	32	16	24	4	8	4	4
AMC	8	16	16	32	8	8	8	16	8	16	16	32	4	8	4	4
LEX	1	4	1	8	1	4	1	1	1	8	1	4	4	8	4	8
CTX	8	16	16	32	32	64	1	16	4	32	1	32	1	32	2	16
IMP	1	4	1	8	1	4	1	1	1	8	1	4	4	8	4	8
GEN	2	8	4	8	4	8	2	8	6	8	2	8	1	8	8	8
AMK	8	8	8	32	16	16	8	16	8	32	8	16	8	32	32	32
CHL	8	16	16	32	8	27	8	16	8	16	8	16	8	8	8	15.2
FOS	32	64	32	64	64	64	32	32	32	32	32	64	32	32	32	32
COL	1	11.6	1	8	1	4	1	7.2	1	8.8	1	23	NA	NA	NA	NA
VAN	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1	2	2	2.4

AMP, ampicillin; AMC, amoxicillin clavulanic acid; LEX, cephalixin; CTX, cefotaxime; IMP, imipenem; GEN, gentamycin; AMK, amikacin; CHL, chloramphenicol; FOS, fosfomycin; COL, colistin; VAN, vancomycin.
NA, not applicable